

ZETA FUNCTIONS OF THE 3-DIMENSIONAL ALMOST BIEBERBACH GROUPS

DIEGO SULCA

ABSTRACT. We present the computations of the subgroup zeta functions of all 3-dimensional almost Bieberbach groups. These computations are an application of the study of zeta functions of virtually nilpotent groups in [S]. We observe that the local factors of these zeta functions satisfy a functional equation as in [V].

1. INTRODUCTION

It has been more than twenty years since Grunewald, Segal and Smith introduced zeta functions of a group in their seminal paper [GSS] as a tool to analyze different sequences associated to it, for example, $\{a_n^{\leq}(G)\}$ where $a_n^{\leq}(G)$ is the number of subgroups of a group G of index n , or $\{a_n^{\triangleleft}(G)\}$ where $a_n^{\triangleleft}(G)$ is the number of normal subgroups of G of index n . The subgroup zeta function of a group G is the Dirichlet series:

$$\zeta_G^{\leq}(s) = \sum_{n=1}^{\infty} \frac{a_n^{\leq}(G)}{n^s} = \sum_{A \leq G} [G : A]^{-s}.$$

When $\zeta_G^{\leq}(s)$ defines an analytic function in some region of $\mathbb{C}$ then interesting properties of $\{a_n^{\leq}(G)\}$ can be obtained such as the degree of its growth and formulas for its asymptotic behavior. The family of groups whose subgroup zeta functions define analytic functions (groups with polynomial subgroup growth) was characterized in [LMS] and it is extremely large including for example all polycyclic-by-finite groups. Unfortunately the family of those groups whose zeta functions is well understood is too small. Most of the analytic properties of subgroup zeta functions have been obtained for the family of finitely generated torsion free nilpotent groups ([GSS],[dSG]). These zeta functions have meromorphic continuation to the left of its abscissa of convergence, which is a rational number, they obey of an Euler product decomposition and the local factors at the prime p are rational functions in p^{-s} .

Date: November 13, 2012.

1991 Mathematics Subject Classification. 20E07, 11M41.

Explicit computations of zeta functions of nilpotent groups are known just for very few groups. Examples are the free abelian groups, some nilpotent groups of Hirsch length ≤ 6 and some other sporadic examples in higher dimension. For a good source of these examples is [dSW]. The situation for non-nilpotent groups is very different and it is remarkable how sensitive the subgroup zeta function is if we extend the group base by a finite group. For example the subgroup zeta function of $\mathbb{Z}$ is just $\zeta(s)$, the Riemann zeta function, which has abscissa of convergence 1, while for the infinite dihedral group D_∞ we have $\zeta_{D_\infty}^{\leq}(s) = \zeta(s) + 2^{-s}\zeta(s-1)$ with abscissa of convergence 2.

A systematic study of zeta functions of virtually abelian groups appears in [dSMS] where it is shown that these zeta functions have meromorphic continuation to the whole plane, that they are a finite sum of series admitting Euler product decomposition and it is given a theoretical expression for these zeta functions in terms of the structure of the group algebra $\mathbb{Q}(G/N)$, where G is the group and N is a finite index normal subgroup isomorphic to $\mathbb{Z}^d$, and the action of G/N on N . There it is also presented the explicit expressions for the zeta functions of the seventeen plane crystallographic groups.

In [S] we study zeta functions of finitely generated virtually nilpotent groups. We express the subgroup zeta function of such a group as a finite sum of Euler product of cone integrals and using results from [dSG] we deduce that these zeta functions have rational abscissa of convergence, that they have meromorphic continuation to the left and we give a method to compute the cone conditions for these integrals. In the current paper we present the computations of the subgroup zeta functions of all 3-dimensional almost Bieberbach groups. It is not difficult to see that this family coincides with the family of all finitely generated torsion free virtually nilpotent groups of Hirsch length 3. This family includes the ten 3-dimensional Bieberbach groups (torsion free space groups) and the others are classified in [DIKL] or [D]. For each 3-dimensional AB -group we shall write a functional equation for the local zeta function which holds for almost all primes.

This article should be read in parallel with [S] because we shall use the same terminology and notation to avoid repetitions. Our computation are checked for the primes $p = 2, 3, 5, 7, 11, 13$ and up to the power p^7 using the Computational Algebra System GAP. We are very grateful with the developers of that software.

2. COMPUTING ZETA FUNCTIONS

An almost Bieberbach group (*AB*-group) is a finitely generated group G containing a finite index normal subgroup N which is nilpotent containing every finite index nilpotent subgroup of G . The subgroup N is the Fitting subgroup of G . The Hirsch length of G is the number of infinite cyclic factors in any subnormal series of G where the factors are finite or cyclic. This number is well defined and it is denoted by $h(G)$ and it is usually called the dimension of the *AB*-group G by geometrical reasons but we shall not go into details. The interested reader should consult [D] for a detailed exposition of these groups.

From now on, this article should be read in parallel with [S] because we shall use the same language and notation. The results in [S] imply that if G is a 3-dimensional *AB*-group with Fitting subgroup N then:

- I. $\zeta_G^{\leq}(s) = \sum_{N \leq H \leq G} [G : H]^{-s} \zeta_{H,N}^{\leq}(s)$ where $\zeta_{H,N}^{\leq}(s) = \sum_{\substack{A \leq_f H \\ AN = H}} [H : A]^{-s}$.
- II. For each $N \leq H \leq G$, we have a decomposition as an Euler product $\zeta_{H,N}^{\leq}(s) = \prod_p \zeta_{H,N,p}^{\leq}(s)$ where $\zeta_{H,N,p}^{\leq}(s) = \sum_{\substack{A \leq_p H \\ AN = H}} [H : A]^{-s}$.
- III. For each $N \leq H \leq G$, there exist finite collections of polynomials $\{f_i\}_{i \in I}$ and $\{g_i\}_{i \in I}$ with rational coefficients such that for each prime p we have

$$\zeta_{H,N,p}^{\leq}(s) = (1 - p^{-1})^{-3} \int_{\mathcal{T}_p} |t_{11}|_p^{s-r} |t_{22}|_p^{s-r-1} |t_{33}|_p^{s-r-2} d\mu$$

where $r = [H : N]$ and

$$\mathcal{T}_p = \{(\mathbf{t}, \mathbf{v}) \in Tr(3, \mathbb{Z}_p) \times M_{r-1 \times 3}(\mathbb{Z}_p) : f_i(\mathbf{t}, \mathbf{v}) | g_i(\mathbf{t}, \mathbf{v}), \forall i \in I\}$$

It is a very important fact that if G is an *AB*-group with Fitting subgroup N and $N \leq H \leq G$, then H is also an *AB*-group with Fitting subgroup N . Then the computation of $\zeta_G^{\leq}(s)$ for all the 3-dimensional *AB*-groups can be obtained from I. above if we know the computation of $\zeta_{G,N}^{\leq}(s)$ for all 3-dimensional *AB*-group G , where N is the Fitting subgroup of G . Hence, we shall only present the computation for $\zeta_{G,N}^{\leq}(s)$.

2.1. 3-dimensional nilpotent AB-groups. The Fitting subgroup of any 3-dimensional *AB*-group is of the form

$$N_k = \langle x_1, x_2, x_3 : [x_2, x_1] = x_3^k, [x_3, x_1] = [x_3, x_2] = 1 \rangle$$

for some non negative integer k . By Proposition 2.6 of [GSS], the local zeta function of N_k at the prime p is given by the integral:

$$\zeta_{N_k,p}^{\leq}(s) = (1 - p^{-1})^{-3} \int_{\mathcal{T}_p} |t_{11}|_p^{s-1} |t_{22}|_p^{s-2} |t_{33}|_p^{s-3} d\mu$$

where $\mathcal{T}_p$ is the set of matrices $\mathbf{t} \in Tr(3, \mathbb{Z}_p)$ representing a good basis for some open subgroup of the pro- p completion of N_k . By Lemma 2.1 of [GSS], a matrix $\mathbf{t}$ represents a good basis if and only if $t_{11}t_{22}t_{33} \neq 0$ and $[\mathbf{x}^{(t_{11}, t_{12}, t_{13})}, \mathbf{x}^{(0, t_{22}, t_{23})}] \in \langle \overline{x_3^{t_{33}}} \rangle$. Doing operations in N_k , the last condition becomes in $x_{33}^{kt_{11}t_{22}} \in \langle \overline{x_3^{t_{33}}} \rangle$, which is clearly equivalent to $t_{33}|kt_{11}t_{22}$. Then we have:

$$\begin{aligned} \zeta_{N_k,p}^{\leq}(s) &= (1 - p^{-1})^{-3} \int_{t_{33}|kt_{11}t_{22}} |t_{11}|_p^{s-1} |t_{22}|_p^{s-2} |t_{33}|_p^{s-3} d\mu \\ &= (1 - p^{-1})^{-3} \sum_{i,j=0}^{\infty} \sum_{l=0}^{v_p(k)+i+j} p^{-i(s-1)} p^{-j(s-2)} p^{-l(s-3)} \int_{\substack{|t_{11}|=p^{-i}, \\ |t_{22}|=p^{-j}, \\ |t_{33}|=p^{-l}}} d\mu \\ &= \sum_{i,j=0}^{\infty} \sum_{l=0}^{v_p(k)+i+j} p^{-is} p^{-j(s-1)} p^{-l(s-2)} \\ &= \zeta_p(s-2) (\zeta_p(s)\zeta_p(s-1) - p^{(2-s)(v_p(k)+1)} \zeta_p(2s-2)\zeta_p(2s-3)). \end{aligned}$$

where v_p is the p -adic valuation on $\mathbb{Z}_p$, and $\zeta_p(s) = \frac{1}{1-p^{-s}}$ is the p -local factor of the Riemann zeta function $\zeta(s)$. Doing the product of these local factors over all the primes p we obtain:

$$\zeta_{N_0}^{\leq}(s) = \zeta(s)\zeta(s-1)\zeta(s-2)$$

which has abscissa of convergence 3; and for $k \neq 0$ we have

$$\begin{aligned} \zeta_{N_k}^{\leq}(s) &= \prod_{p|k} \frac{\zeta_p(s)\zeta_p(s-1) - p^{(2-s)(v_p(k)+1)} \zeta_p(2s-2)\zeta_p(2s-3)}{\zeta_p(s)\zeta_p(s-1) - p^{(2-s)} \zeta_p(2s-2)\zeta_p(2s-3)} \\ &\quad \cdot \frac{\zeta(s)\zeta(s-1)\zeta(2s-2)\zeta(2s-3)}{\zeta(3s-3)} \end{aligned}$$

and Corollary 1.10 of [S] shows that the abscissa of convergence is 2. This concludes the calculation of the zeta functions all 3-dimensional nilpotent AB-groups. In fact, $\{N_k : k \geq 0\}$ are the representatives of the different isomorphism classes.

2.2. 3-dimensional non-nilpotent AB-groups. When G is a 3-dimensional non-nilpotent AB-group with Fitting subgroup N , to compute $\zeta_{G,N}^{\leq}(s)$ we shall follow the following steps:

Step 1. We shall present the 3-dimensional AB -group G with generators $x_1, x_2, x_3, \alpha, \beta, \dots$ and a set of relations, such that $\{x_1, x_2, x_3\}$ is a Mal'cev basis for the Fitting subgroup, say $N = N_k$.

Step 2. A simplified expression for $\zeta_{G,N,p}^{\leq}(s)$ as an integral is given taking advantage of the presentation of the group G/N :

Proposition 2.1. *Suppose that the group G/N is defined by t generators $\gamma_{i_1}N, \dots, \gamma_{i_t}N$ and a finite set of relations $R_j(\gamma_{i_1}, \dots, \gamma_{i_t}) \in N$ ($j \in J$). Then for each prime p we have*

$$(2.1) \quad \zeta_{G,N,p}^{\leq}(s) = (1 - p^{-1})^{-3} \int_{\mathcal{T}'_p} |t_{11}|^{s-1-t} |t_{22}|^{s-2-t} |t_{33}|^{s-3-t} d\mu$$

where $\mathcal{T}'_p$ is the set of pairs $(\mathbf{t}, \mathbf{v}) \in \text{Tr}(3, \mathbb{Z}_p) \times M_{r-1 \times 3}(\mathbb{Z}_p)$ satisfying the following conditions:

- (1) $t_{33} | kt_{11}t_{22}$
- (2) $(\gamma_{i_j} \mathbf{x}^{\mathbf{v}_{i_j}})^{-1} \mathbf{x}^{\mathbf{t}_{i_j}} \gamma_{i_j} \mathbf{x}^{\mathbf{v}_{i_j}} \in B_{\mathbf{t}}, \quad i = 1, \dots, h, \quad j = 1, \dots, t$
- (3) $R_j(\gamma_{i_1} \mathbf{x}^{\mathbf{v}_{i_1}}, \dots, \gamma_{i_t} \mathbf{x}^{\mathbf{v}_{i_t}}) \in B_{\mathbf{t}}, \quad j \in J$.

Here and in the proof below the notation is $B_{\mathbf{t}} = \overline{\langle \mathbf{x}^{\mathbf{t}_1}, \mathbf{x}^{\mathbf{t}_2}, \mathbf{x}^{\mathbf{t}_3} \rangle}$ and $A_{(\mathbf{t}, \mathbf{v})} = B_{\mathbf{t}} \cup (\cup_{i=1}^{r-1} \gamma_i \mathbf{x}^{\mathbf{v}_i} B_{\mathbf{t}})$.

Proof. We shall follow the proof of Proposition 1.8 in [S]. For $j \notin \{i_1, \dots, i_t\}$ there exists a word w_j such that $\gamma_j N = w_j(\gamma_{i_1}, \dots, \gamma_{i_t})N$. We claim that $\mathcal{T}_p$ is the same as the set of pairs $(\mathbf{t}, \mathbf{v})$ such that

- (1) $t_{33} | kt_{11}t_{22}$
- (2) $(\gamma_j \mathbf{x}^{\mathbf{v}_j})^{-1} \mathbf{x}^{\mathbf{t}_j} \gamma_j \mathbf{x}^{\mathbf{v}_j} \in B_{\mathbf{t}}, \quad i = 1, \dots, h, \quad j = 1, \dots, r-1$
- (3) $R_j(\gamma_{i_1} \mathbf{x}^{\mathbf{v}_{i_1}}, \dots, \gamma_{i_t} \mathbf{x}^{\mathbf{v}_{i_t}}) \in B_{\mathbf{t}}, \quad j \in J$.
- (4) $\gamma_j \mathbf{x}^{\mathbf{v}_j} \in w_j(\gamma_{i_1} \mathbf{x}^{\mathbf{v}_{i_1}}, \dots, \gamma_{i_t} \mathbf{x}^{\mathbf{v}_{i_t}}) B_{\mathbf{t}}$ for $j \notin B\{i_1, \dots, i_t\}$.

In fact, any $(\mathbf{t}, \mathbf{v}) \in \mathcal{T}_p$ satisfies (1) and (2). Condition (3) follows since $R_j(\gamma_{i_1} \mathbf{x}^{\mathbf{v}_{i_1}}, \dots, \gamma_{i_t} \mathbf{x}^{\mathbf{v}_{i_t}}) \in A_{(\mathbf{t}, \mathbf{v})} \cap N_p = B_{\mathbf{t}}$, and for (4) we have $(\gamma_j \mathbf{x}^{\mathbf{v}_j})^{-1} w_j(\gamma_{i_1} \mathbf{x}^{\mathbf{v}_{i_1}}, \dots, \gamma_{i_t} \mathbf{x}^{\mathbf{v}_{i_t}}) \in A_{(\mathbf{t}, \mathbf{v})} \cap N = B_{\mathbf{t}}$. For the converse, suppose that a pair $(\mathbf{t}, \mathbf{v})$ satisfies (1)-(4). Condition (1) says that $B_{\mathbf{t}}$ is a subgroup and (2) says that the elements $\gamma_i \mathbf{x}^{\mathbf{v}_i} B_{\mathbf{t}}$ are in $N_{G_p}(B_{\mathbf{t}})/B_{\mathbf{t}}$. Let $A/B_{\mathbf{t}}$ be the subgroup of $N_{G_p}(B_{\mathbf{t}})/B_{\mathbf{t}}$ generated by $\{\gamma_{i_1} \mathbf{x}^{\mathbf{v}_{i_1}} B_{\mathbf{t}}, \dots, \gamma_{i_t} \mathbf{x}^{\mathbf{v}_{i_t}} B_{\mathbf{t}}\}$. Condition (4) says that every $\gamma_i \mathbf{x}^{\mathbf{v}_i} B_{\mathbf{t}}$ is in $A/B_{\mathbf{t}}$ and condition (3) says that $A/B_{\mathbf{t}}$ satisfies the relations R_j ($j \in J$). Then $A/B_{\mathbf{t}}$ is a quotient of G/N and therefore it must be of order $\leq r$. Since $A_{(\mathbf{t}, \mathbf{v})}$ has r elements then $A_{(\mathbf{t}, \mathbf{v})} = A$ and it is a group.

Now observe that once condition (4) is verified we only need to verify condition (2) for those $j \in \{i_1, \dots, i_t\}$. Let's refer to this new condition as (2'). Then when computing the integral, we can start integrating

on the variables v_{jl} for those $j \notin \{i_1, \dots, i_t\}$, that is, on condition (4). Using Proposition 1.3 of [S] we obtain that this partial result is $[N_k : B_{\mathbf{t}}]^{r-1-t} = |t_{11}|^{r-1-t}|t_{22}|^{r-1-t}|t_{33}|^{r-1-t}$. Finally we have to integrate $|t_{11}|^{s-1-t}|t_{22}|^{s-2-t}|t_{33}|^{s-3-t}$ on the set of pairs $(\mathbf{t}, \mathbf{v})$ satisfying conditions (1), (2') and (3), and this is exactly the statement of the proposition. $\square$

Step 3. The conditions which describe the domain of integration in Proposition 2.1 are presented as a set of conditions of the form $\mathbf{x}^{\mathbf{h}(\mathbf{t}, \mathbf{v})} \in B_{\mathbf{t}}$ for some vectorial polynomials $\mathbf{h}$ which are independent of p . This can be done using the definition of G as in the proof of Proposition 1.8 in [S].

Step 4. We present the cone conditions associated to the conditions obtained in *Step 3*. To do this we shall use Proposition 1.5 of [S] which, applied to N_k , gives: If $\mathbf{t}$ represents a good basis for some open subgroup $B_{\mathbf{t}}$ of the pro- p completion of $N = N_k$, then an element $\mathbf{x}^{(a_1, a_2, a_3)}$ is in $B_{\mathbf{t}}$ if and only if the following is true:

$$\begin{aligned} (1) & \quad t_{11} | a_1 \\ (2) & \quad t_{22} | -\frac{a_1}{t_{11}} t_{12} + a_2 \\ (3) & \quad t_{33} | -\frac{-\frac{a_1}{t_{11}} + a_2}{t_{22}} t_{23} - \frac{a_1}{t_{11}} t_{13} + k \frac{a_1}{t_{11}} (\frac{a_1}{t_{11}} + 1) t_{11} t_{12} - k a_1 a_2 + a_3 \end{aligned}$$

Step 5. We start the process of simplification of our set of conditions. First we eliminate those conditions which are implied by the others or at least we shall simplify conditions using the others, for example, $a|b+c$ and $a|c$ will be changed by $a|b$ and $a|c$. Sometimes it will be convenient to distinguish between the two cases $p|[G : N]$ and $p \nmid [G : N]$. For example, if $p \nmid [G : N]$ then any divisor of $[G : N]$ will be a unit and a condition of the form $a|[G : N]b$ is just $a|b$, whereas if $p|[G : N]$ then a condition of the form $a|[G : N]b + 1$ says that a is a unit and therefore we can replace this condition by $a \in \mathbb{Z}_p^*$. Sometimes it will be necessary to split the domain of integration, that is, consider cases: for example the condition $a|b$ can be treated in two cases: the case $|a| = |b|$ and the case $|a| > |b|$. We compute the integral in the two cases and then we sum the results of them. The aim of these simplifications should be clear from the following remark:

Remark 2.2. Let $f_0, g_0 \in \mathbb{Z}_p[x_1, \dots, x_m]$, D_0 a measurable subset of $\mathbb{Z}_p^m$ and for $i = 1, \dots, n$ let D_i be a subset of $\mathbb{Z}_p^m \times \mathbb{Z}_p^i$. Suppose that there exist monomials $g_1, \dots, g_n \in \mathbb{Q}_p[x_1, \dots, x_m]$ and rational functions $k_i \in \mathbb{Q}_p(x_1, \dots, x_n, y_1, \dots, y_{i-1})$ for $i = 1, \dots, n$ such that $g_i(D_0) \subseteq \mathbb{Z}_p$, $k_i(D_{i-1}) \subseteq \mathbb{Z}_p$ and such that $D_i = D_{i-1} \times V_i$, where V_i is

a subset of $\mathbb{Z}_p$ of one of the following two types:

- I: $V_i = \{v_i \in \mathbb{Z}_p : g_i(t_1, \dots, t_m) | k_i(t_1, \dots, t_m, v_1, \dots, v_{i-1}) + v_i\}$
 II: $V_i = \{v_i \in \mathbb{Z}_p : |g_i(t_1, \dots, t_m)| = |k_i(t_1, \dots, t_m, v_1, \dots, v_{i-1}) + \lambda v_i|\}$

Then

$$\int_{D_n} |f_0(t_1, \dots, t_m)|^s |g_0(t_1, \dots, t_m)| d\mu = (1 - p^{-1})^r \int_{D_0} |f_0(t_1, \dots, t_m)|^s \prod_{i=0}^n |g_i(t_1, \dots, t_m)| d\mu$$

where r is the number of sets V_i of type II. In fact this will be an application of Fubini's theorem. We start integrating in the variable v_n and we observe that $g_n(t_1, \dots, t_m) | k_n(t_1, \dots, t_m, v_1, \dots, v_{i-1}) + v_i$ if and only if $v_i \in g_n(t_1, \dots, t_m) \mathbb{Z}_p - k_n(t_1, \dots, t_m, v_1, \dots, v_{i-1})$, and the Haar measure of $g_n(t_1, \dots, t_m) \mathbb{Z}_p - k_n(t_1, \dots, t_m, v_1, \dots, v_{i-1})$ is just $|g_n(t_1, \dots, t_m)|$; whereas $|g_i(t_1, \dots, t_m)| = |k_i(t_1, \dots, t_m, v_1, \dots, v_{i-1}) + v_i|$ if and only if $v_i \in g_i(t_1, \dots, t_m) \mathbb{Z}_p^* - k_i(t_1, \dots, t_m, v_1, \dots, v_{i-1})$ and the Haar measure of $g_i(t_1, \dots, t_m) \mathbb{Z}_p^* - k_i(t_1, \dots, t_m, v_1, \dots, v_{i-1})$ is $(1 - p^{-1}) |g_n(t_1, \dots, t_m)|$. Then integrating on v_n is the same as adding the factor $|g_n(t_1, \dots, t_m)|$ or $(1 - p^{-1}) |g_n(t_1, \dots, t_m)|$ to the integrand, according to when V_n is of the type I or II. Then we continue with the variable v_{n-1} and so on. For example, if $D_0 = \mathbb{Z}_p^m$ and all V_i 's are of type I, then D_n is the subset of $\mathbb{Z}_p^m \times \mathbb{Z}_p^n$ such that the following holds:

$$\begin{aligned} & g_1(t_1, \dots, t_m) | k_1(t_1, \dots, t_m) + v_1; \\ & g_2(t_1, \dots, t_m) | k_2(t_1, \dots, t_m, v_1) + v_2; \\ & \vdots \\ & g_n(t_1, \dots, t_m) | k_n(t_1, \dots, t_m, v_1, \dots, v_{n-1}) + v_n. \end{aligned}$$

We shall use this remark where t_1, t_2, t_3 will be the variables t_{11}, t_{22} and t_{33} and $v_1, \dots, v_n$ will be the other variables.

Step 6: After computing all the local factors we shall do the Euler product of them and we will write down the explicit formula for $\zeta_{G,N}^{\leq}(s)$. We will be able to read the abscissa of convergence from this expression according to Corollary 1.10 of [S]. Most of these expressions can be written in terms of the Riemann zeta function $\zeta(s) = \prod_p \zeta_p(s) = \prod_p \frac{1}{1-p^{-s}}$ and sometimes in terms of Dirichlet L-series $L(s, \chi) = \prod_p L(s, \chi, p) = \prod_p \frac{1}{1-\chi(p)p^{-s}}$ of some extended primitive residue class character $\chi : \mathbb{Z} \rightarrow (\mathbb{Z}/n\mathbb{Z})^*$. Those which will appear

are χ_3 and χ_4 defined by:

$$\chi_3(a) = \begin{cases} 1 & \text{if } a \equiv 1 \pmod{3}, \\ -1 & \text{if } a \equiv 2 \pmod{3}, \\ 0 & \text{otherwise} \end{cases} \text{ and } \chi_4(a) = \begin{cases} 1 & \text{if } a \equiv 1 \pmod{4}, \\ -1 & \text{if } a \equiv 3 \pmod{4}, \\ 0 & \text{otherwise} \end{cases}$$

It is a fact that any Dirichlet L -series, $L(s, \chi)$, has abscissa of convergence 1.

Step 7: In [V] it is proven that for a τ -group N , the local zeta function $\zeta_{N,p}^{\leq}(s)$ satisfies a functional equation $\zeta_{N,p}^{\leq}(s)|_{p \rightarrow p^{-1}} = p^{-ns + \binom{n}{2}} \zeta_{N,p}^{\leq}(s)$ for almost all primes p . We shall present functional equations for the local zeta functions $\zeta_{G,N,p}^{\leq}(s)$ for almost all primes p .

We begin with the nine 3-dimensional Bieberbach (the abelian case was treated before) giving its IT-number in the International Table for Crystallography. Then we continue with the AB -groups given as in [D].

2.2.1. *The Bieberbach group $\mathfrak{G}_2$: $IT=(3,4)$.* This is the group

$$\mathfrak{G}_2 = \langle \gamma, x_1, x_3, x_3 : [x_i, x_j] = 1, \gamma^2 = x_1, \gamma x_2 \gamma^{-1} = x_2^{-1}, \gamma x_3 \gamma^{-1} = x_3^{-1} \rangle$$

with Fitting subgroup $N_0 = \langle x_1, x_2, x_3 \rangle$. We have

$$(2.2) \quad \zeta_{\mathfrak{G}_2, N_0, p}^{\leq}(s) = (1 - p^{-1})^{-3} \int_{\mathcal{T}'} |t_{11}|^{s-2} |t_{22}|^{s-3} |t_{33}|^{s-4} d\mu$$

where $\mathcal{T}'$ is the set of pairs $(\mathbf{t}, \mathbf{v}) \in Tr(3, \mathbb{Z}_p) \times \mathbb{Z}_p^3$ for which

$$\gamma \mathbf{x}^{\mathbf{v}} \mathbf{x}^{\mathbf{t}_i} (\gamma \mathbf{x}^{\mathbf{v}})^{-1} \in B_{\mathbf{t}} \text{ for } i = 1, 2, 3 \text{ and } (\gamma \mathbf{x}^{\mathbf{v}})^2 \in B_{\mathbf{t}}.$$

These conditions are translated in:

$$x_2^{2t_{12}} x_3^{2t_{13}} \in B_{\mathbf{t}}, \text{ and } x_1^{2v_1+1} \in B_{\mathbf{t}},$$

whose associated cone conditions are:

- (1) $t_{22} | 2t_{12},$
- (2) $t_{33} | -\frac{2t_{12}}{t_{22}} t_{23} + 2t_{13},$
- (3) $t_{11} | 2v_1 + 1,$
- (4) $t_{22} | -\frac{2v_1+1}{t_{11}} t_{12}$
- (5) $t_{33} | \frac{2v_1+1}{t_{11}} \frac{t_{12}}{t_{22}} t_{23} - \frac{2v_1+1}{t_{11}} t_{23}.$

When $p \neq 2$ then (1) implies (4) and (2) implies (5) and therefore the reduced conditions are:

$$t_{22} | t_{12}; t_{33} | -\frac{t_{12}t_{23}}{t_{22}} + t_{13} \text{ and } t_{11} | v_1 + 2^{-1}.$$

Using Remark 2.2 with $D_0 = \mathbb{Z}_p^3$, we transform the integral (2.2) into

$$\zeta_{\mathfrak{G}_2, N_0, p}^{\leq}(s) = (1-p^{-1})^{-3} \int |t_{11}|^{s-1} |t_{22}|^{s-2} |t_{33}|^{s-3} = \zeta_p(s) \zeta_p(s-1) \zeta_p(s-2).$$

When $p = 2$, then condition (3) can be changed by $t_{11} \in \mathbb{Z}_2^*$, (4) by $t_{22}|t_{12}$ which implies (1), and (5) by $t_{33}|\frac{t_{12}t_{23}}{t_{22}} - t_{23}$ which implies (2). Then our conditions are

$$t_{11} \in \mathbb{Z}_2^*, \quad t_{22}|t_{12}, \quad \text{and} \quad t_{33}|\frac{t_{12}t_{23}}{t_{22}} - t_{23}.$$

Applying Remark 2.2 with $D_0 = \mathbb{Z}_2^* \times \mathbb{Z}_2^2$ we transform the integral (2.2) in

$$\zeta_{\mathfrak{G}_2, N_0, 2}^{\leq}(s) = (1-2^{-1})^{-3} \int_{t_{11} \in \mathbb{Z}_2^*} |t_{22}|^{s-2} |t_{33}|^{s-3} d\mu = \zeta_2(s-1) \zeta_2(s-2).$$

Doing the Euler product of all these local factors we obtain:

$$(2.3) \quad \zeta_{\mathfrak{G}_2, N}^{\leq}(s) = \zeta(s) \zeta(s-1) \zeta(s-2) \zeta_2(s)^{-1},$$

which has abscissa of convergence 3. The functional equation for almost all primes is:

$$(2.4) \quad \zeta_{\mathfrak{G}_2, N, p}^{\leq}(s)|_{p \rightarrow p^{-1}} = (-1)^3 p^{-3s+3} \zeta_{\mathfrak{G}_2, N, p}^{\leq}(s).$$

2.2.2. The Bieberbach group $\mathfrak{G}_3$, $IT=(3,144)$. This is the group

$$\mathfrak{G}_3 = \langle \gamma, x_1, x_2, x_3 : [x_i, x_j] = 1, \gamma^3 = x_1, \gamma x_2 \gamma^{-1} = x_3, \gamma x_3 \gamma^{-1} = x_2^{-1} x_3^{-1} \rangle$$

with Fitting subgroup $N_0 = \langle x_1, x_2, x_3 \rangle$. We have

$$(2.5) \quad \zeta_{\mathfrak{G}_3, N_0, p}^{\leq}(s) = (1-p^{-1})^{-3} \int_{\mathcal{T}'} |t_{11}|^{s-2} |t_{22}|^{s-3} |t_{33}|^{s-4} d\mu$$

where $\mathcal{T}'$ is the set of all pairs $(\mathbf{t}, \mathbf{v}) \in Tr(3, \mathbb{Z}_p) \times \mathbb{Z}_p^3$ satisfying:

$$(\gamma \mathbf{x}^{\mathbf{v}}) \mathbf{x}^{\mathbf{t}_i} (\gamma \mathbf{x}^{\mathbf{v}})^{-1} \in B_{\mathbf{t}} \text{ for } i = 1, 2, 3; \text{ and } (\gamma \mathbf{x}^{\mathbf{v}})^3 \in B_{\mathbf{t}}.$$

These conditions are translated in

$$x_2^{t_{13}+t_{12}} x_3^{2t_{13}-t_{12}} \in B_{\mathbf{t}}, \quad x_2^{t_{23}} x_3^{t_{23}-t_{22}} \in B_{\mathbf{t}}, \quad x_2^{t_{33}} \in B_{\mathbf{t}}, \quad x_1^{3v_1+1} \in B_{\mathbf{t}}.$$

The associated cone conditions are:

- (1) $t_{22}|t_{13} + t_{12};$
- (2) $t_{33} | -\frac{t_{13}+t_{12}}{t_{22}} t_{23} + 2t_{13} - t_{12};$
- (3) $t_{22}|t_{23};$
- (4) $t_{33} | -\frac{t_{23}}{t_{22}} t_{23} + t_{23} - t_{22};$
- (5) $t_{22}|t_{33};$
- (6) $t_{33} | -\frac{t_{33}}{t_{22}} t_{23};$
- (7) $t_{11}|3v_1 + 1;$

- (8) $t_{22} \mid -\frac{3v_1+1}{t_{11}}t_{12}$;
- (9) $t_{33} \mid \frac{3v_1+1}{t_{11}}(\frac{t_{12}}{t_{22}}t_{23} - t_{13})$;

Condition (3) implies (6) and conditions (4) and (5) imply $t_{22} \mid -\frac{t_{23}}{t_{22}}t_{23} + t_{23} - t_{22}$, or equivalently $(\frac{t_{23}}{t_{22}})^2 - \frac{t_{23}}{t_{22}} \in \mathbb{Z}_p$, which is clearly possible only if $t_{22} \mid t_{23}$. Then (4) and (5) imply (3). Let's see that (1) and (8) can be changed by $t_{22} \mid t_{12}$ and $t_{22} \mid t_{13}$. In fact, it is clear that this new condition implies (1) and (8), and conversely if $p = 3$ then (8) is the same as $t_{22} \mid t_{12}$ which, in combination with (1), implies also $t_{22} \mid t_{13}$, whereas if $p \neq 3$, then (5) and (2) imply $t_{22} \mid 2t_{13} - t_{12}$, and this in combination with (1) implies $t_{22} \mid 3t_{13}$, which is the same as $t_{22} \mid t_{13}$ and with (1) this implies $t_{22} \mid t_{12}$.

Now we shall see that conditions (2) and (9) can be replaced by a new condition: (*) $t_{33} \mid \frac{t_{12}}{t_{22}}t_{23} - t_{13}$. It is enough to see that (*) can be deduced from (2) and (9) because (*) clearly implies (9) and multiplying (*) by $\frac{t_{23}}{t_{22}}$ we obtain $t_{33} \mid \frac{t_{12}}{t_{22}}\frac{t_{23}^2}{t_{22}} - \frac{t_{13}t_{23}}{t_{22}}$, which by condition (4), is equivalent to $t_{33} \mid \frac{t_{12}}{t_{22}}(t_{23} - t_{22}) - \frac{t_{13}t_{23}}{t_{22}}$, and therefore we have $t_{33} \mid \frac{t_{12}}{t_{22}}(t_{23} - t_{22}) - \frac{t_{13}t_{23}}{t_{22}} - 2(\frac{t_{12}}{t_{22}} - t_{13}) = -\frac{t_{13}+t_{12}}{t_{22}}t_{23} + 2t_{13} - t_{12}$, which is condition (2). Now observe that if $p = 3$ then (*) follows from (9). If $p \neq 3$ then multiplying the right hand side of (2) by $\frac{t_{23}}{t_{22}} + 1$ we obtain $t_{33} \mid (-\frac{t_{13}+t_{12}}{t_{22}}t_{23} + 2t_{13} - t_{12})(\frac{t_{23}}{t_{22}} + 1) = -t_{12}((\frac{t_{23}}{t_{22}})^2 + 2\frac{t_{23}}{t_{22}} + 1) + t_{13}(-(\frac{t_{23}}{t_{22}})^2 + \frac{t_{23}}{t_{22}} + 2)$ which, by condition (4), can be reduced into $t_{33} \mid -3\frac{t_{12}}{t_{22}}t_{23} + 3t_{13}$ and this implies (*).

Finally (*), (4) and (5) clearly imply $t_{22} \mid t_{13}$ and our final set of conditions is:

- (1') $t_{22} \mid t_{33}$
- (2') $t_{22} \mid t_{12}$;
- (3') $t_{22}t_{33} \mid -t_{23}^2 + t_{23}t_{22} - t_{22}^2$;
- (4') $t_{11} \mid 3v_1 + 1$ (or $t_{11} \in \mathbb{Z}_3^*$ if $p = 3$);
- (5') $t_{33} \mid \frac{t_{12}}{t_{22}}t_{23} - t_{13}$.

Suppose that $p = 3$. We claim that condition (3') holds if and only if $|t_{22}| = |t_{33}|$ and $t_{22} \mid t_{23}$ or $|t_{22}| = 3|t_{33}|$ and $3t_{22} \mid t_{23} - 3t_{22}$. In fact, if $|t_{22}| = |t_{33}|$ then (3') is reduced to $t_{22}^2 \mid t_{23}(t_{23} - t_{22})$ which is equivalent to $t_{22} \mid t_{23}$. If $|t_{22}| = 3|t_{33}|$ then (3') is the same as $3 \mid -(\frac{t_{23}}{t_{22}})^2 + \frac{t_{23}}{t_{22}} - 1$ and the last condition happens if and only if $\frac{t_{23}}{t_{22}} \in 2 + 3\mathbb{Z}_3$, that is, $3t_{22} \mid t_{23} - 2t_{22}$. Finally if $3^2t_{22} \mid t_{33}$ then (3') would imply $9 \mid -(\frac{t_{23}}{t_{22}})^2 + \frac{t_{23}}{t_{22}} - 1$ but there is not solution of $x^2 - x + 1 = 0 \pmod{9}$. Applying Remark 2.2 in both

cases we obtain:

$$(1 - 3^{-1})^{-3} \int_{t_{11} \in \mathbb{Z}_3^*, |t_{22}|=|t_{33}|} |t_{22}|^{s-1} |t_{33}|^{s-3} d\mu = \zeta_3(2s-2),$$

and

$$(1 - 3^{-1})^{-3} \int_{t_{11} \in \mathbb{Z}_3^*, |t_{22}|=3|t_{33}|} 3^{-1} |t_{22}|^{s-1} |t_{33}|^{s-3} d\mu = 3^{1-s} \zeta_3(2s-2)$$

Then we obtain

$$\zeta_{\mathfrak{G}_3, N_0, 3}^{\leq}(s) = (1 + 3^{1-s}) \zeta_3(2s-2) = \zeta_3(s-1).$$

Suppose that $p \neq 3$. When $|t_{22}| = |t_{33}|$ then (3') is equivalent to $t_{22}|t_{23}$ and we can apply Remark 2.2 to obtain

$$(1 - p^{-1})^{-3} \int_{|t_{22}|=|t_{33}|} |t_{11}|^{s-1} |t_{22}|^{s-1} |t_{33}|^{s-3} d\mu = \zeta_p(s) \zeta_p(2s-2)$$

If $|t_{22}| > |t_{33}|$ then (3') implies that $\frac{t_{23}}{t_{22}}$ is a solution of $x^2 - x + 1 = 0 \pmod{p}$ which can happen only if $p \equiv 1 \pmod{3}$ and so we assume that this is the case. By Hensel's lemma there exist $\eta_1, \eta_2 \in \mathbb{Z}_p$ such that $x^2 - x + 1 = (x - \eta_1)(x - \eta_2)$. Then it is easy to see that condition (3') becomes equivalent to $t_{33}| - (t_{23} - \eta_1 t_{22})(t_{23} - \eta_2 t_{22})$ and this is equivalent to the following condition

$$t_{33}|t_{23} - \eta_1 t_{22} \text{ or } t_{33}|t_{23} - \eta_2 t_{22}$$

It is easy to see that these two cases split our domain in two disjoint subsets and therefore we can apply Remark 2.2 in both cases. Their sum is

$$2(1 - p^{-1})^{-3} \int_{|t_{22}| > |t_{33}|} |t_{11}|^{s-1} |t_{22}|^{s-2} |t_{33}|^{s-2} d\mu = 2p^{1-s} \zeta_p(s) \zeta_p(s-1) \zeta_p(2s-2).$$

To write down the final result we shall use the Dirichlet character χ_3 to obtain:

$$\begin{aligned} \zeta_{\mathfrak{G}_3, N_0, p}^{\leq}(s) &= \zeta_p(s) \zeta_p(2s-2) \left(1 + (\chi_3(p) + 1) \frac{p^{1-s}}{1 - p^{1-s}} \right) \\ &= \zeta_p(s) \zeta_p(2s-2) \frac{1 + \chi_3(p) p^{1-s}}{1 - p^{1-s}} \\ &= \zeta_p(s) \zeta_p(2s-2) \frac{1 - p^{2-2s}}{1 - p^{1-s}} \frac{1}{1 - \chi_3(p) p^{1-s}} \\ &= \zeta_p(s) \zeta_p(s-1) L(s-1, \chi_3, p) \end{aligned}$$

The Euler product of all these local factors is:

$$(2.6) \quad \zeta_{\mathfrak{G}_3, N_0}^{\leq}(s) = \zeta(s)\zeta(s-1)L(s-1, \chi_3) \frac{1}{\zeta_3(s)L(s-1, \chi_3, 3)}$$

which has abscissa of convergence 2. The functional equation for almost all primes p is:

$$(2.7) \quad \zeta_{\mathfrak{G}_3, N_0, p}^{\leq}(s)|_{p \rightarrow p^{-1}} = (-1)^3 \chi_3(p) p^{-3s+2} \zeta_{\mathfrak{G}_3, N_0, p}^{\leq}(s).$$

2.2.3. *The Bieberbach group $\mathfrak{G}_4$: $IT=(3, 76)$.* This is the group

$$\mathfrak{G}_4 = \langle \gamma, x_1, x_2, x_3 : [x_i, x_j] = 1, \gamma^4 = x_1, \gamma x_2 \gamma^{-1} = x_3, \gamma x_3 \gamma^{-1} = x_2^{-1} \rangle$$

with Fitting subgroup $N_0 = \langle x_1, x_2, x_3 \rangle$. We have

$$(2.8) \quad \zeta_{\mathfrak{G}_4, N_0, p}^{\leq}(s) = (1 - p^{-1})^{-3} \int_{\mathcal{T}'} |t_{11}|^{s-2} |t_{22}|^{s-3} |t_{33}|^{s-4} d\mu$$

where $\mathcal{T}'$ is the set of pairs $(\mathbf{t}, \mathbf{v}) \in Tr(3, \mathbb{Z}_p) \times \mathbb{Z}_p^3$ satisfying

$$\gamma \mathbf{x}^{\mathbf{v}} \mathbf{x}^{\mathbf{t}_i} (\gamma \mathbf{x}^{\mathbf{v}})^{-1} \in B_{\mathbf{t}} \text{ for } i = 1, 2, 3 \text{ and } (\gamma \mathbf{x}^{\mathbf{v}})^4 \in B_{\mathbf{t}}.$$

These conditions are translated in:

$$x_2^{t_{12}+t_{13}} x_3^{-t_{12}+t_{13}} \in B_{\mathbf{t}}, \quad x_2^{-t_{23}} x_3^{t_{22}} \in B_{\mathbf{t}}, \quad x_2^{t_{33}} \in B_{\mathbf{t}}, \quad x_1^{4v_1+1} \in B_{\mathbf{t}}.$$

whose associated cone conditions are:

- (1) $t_{22}|t_{12} + t_{13}$;
- (2) $t_{33} | -\frac{t_{12}+t_{13}}{t_{22}} t_{23} - t_{12} + t_{13}$;
- (3) $t_{22}|t_{23}$;
- (4) $t_{33} | \frac{t_{23}}{t_{22}} t_{23} + t_{22}$;
- (5) $t_{22}|t_{33}$;
- (6) $t_{11}|4v_1 + 1$;
- (7) $t_{22} | -\frac{4v_1+1}{t_{11}} t_{12}$;
- (8) $t_{33} | \frac{4v_1+1}{t_{11}} (\frac{t_{12}}{t_{22}} t_{23} - t_{13})$.

Conditions (1) and (7) can be replaced by $t_{22}|t_{12}$ and $t_{22}|t_{13}$. In fact, for $p = 2$ this follows easily using (7) while for $p \neq 2$ we use conditions (2) and (3) in combination with (1) to obtain $t_{22}|2t_{12}$ which implies $t_{22}|t_{12}$. Similarly (4) and (5) imply $t_{22}^2|t_{23}^2 + t_{22}^2$ which implies (3).

Now we shall see that conditions (2) and (8) can be replaced by (*) $t_{33} | \frac{t_{12}}{t_{22}} t_{23} - t_{13}$. It is enough to see that we can deduce (*) from the others

because (*) imply (8) and multiplying (*) by $\frac{t_{23}}{t_{22}}$ we obtain $t_{33} | \frac{t_{12}}{t_{22}} \frac{t_{23}^2}{t_{22}} - t_{13} \frac{t_{23}}{t_{22}}$, which in combination with (4) gives $t_{33} | t_{12} - t_{13} \frac{t_{23}}{t_{22}}$. Then $t_{33} | t_{12} - t_{13} \frac{t_{23}}{t_{22}} + \frac{t_{12}}{t_{22}} t_{23} - t_{13}$ which is equivalent to (3). Now, from (3) we have $t_{33} | (-\frac{t_{12}+t_{13}}{t_{22}} t_{23} - t_{12} + t_{13})(\frac{t_{23}}{t_{22}} + 1)$. Straightforward computations on the right hand side of this condition, replacing $\frac{t_{23}}{t_{22}}$ by $-t_{22}$ which is

possible by (4), shows that $t_{33} \mid -2\frac{t_{12}}{t_{22}}t_{23} + 2t_{13}$. If $p \neq 2$ then this implies (*) and if $p = 2$ then (*) is implied by (8).

Finally observe that (*) in combination with (5) implies $t_{22} \mid t_{13}$. Then our new set of conditions is

- (1') $t_{22} \mid t_{33}$;
- (2') $t_{22} \mid t_{12}$;
- (3') $t_{22}t_{33} \mid t_{23}^2 + t_{22}^2$;
- (4') $t_{11} \mid 4v_1 + 1$ (or $t_{11} \in \mathbb{Z}_2^*$ if $p = 2$);
- (5') $t_{33} \mid \frac{t_{12}}{t_{22}}t_{23} - t_{13}$.

When $p = 2$ then it is easy to see, as in the last case, that (3') holds if and only if $|t_{22}| = |t_{33}|$ and $t_{22} \mid t_{23}$, or $|t_{22}| = 2|t_{33}|$ and $2t_{22} \mid t_{23} - t_{22}$ (here we use that $x^2 + 1 = 0$ has no solution mod 4). Then we compute the integral in both cases according to Remark 2.2 and the results are respectively

$$(1 - 2^{-1})^{-3} \int_{t_{11} \in \mathbb{Z}_2^*, |t_{22}| = |t_{33}|} |t_{22}|^{s-1} |t_{33}|^{s-3} d\mu = \zeta_2(2s - 2),$$

and

$$(1 - 2^{-1})^{-3} \int_{t_{11} \in \mathbb{Z}_2^*, |t_{22}| = 2|t_{33}|} 2^{-1} |t_{22}|^{s-1} |t_{33}|^{s-3} d\mu = 2^{1-s} \zeta_2(2s - 2)$$

and therefore

$$\zeta_{\mathfrak{G}_{4,N_0,2}}^{\leq}(s) = (1 + 2^{1-s}) \zeta_2(2s - 2) = \zeta_2(s - 1).$$

Suppose that $p \neq 2$. If $|t_{22}| = |t_{33}|$ then condition (3') is equivalent to $t_{22} \mid t_{23}$ and the integral in this case is

$$(1 - p^{-1})^{-3} \int_{|t_{22}| = |t_{33}|} |t_{11}|^{s-1} |t_{22}|^{s-1} |t_{33}|^{s-3} d\mu = \zeta_p(s) \zeta_p(2s - 2)$$

If $|t_{22}| > |t_{33}|$ then condition (3') is $t_{33} \mid t_{22}((\frac{t_{23}}{t_{22}})^2 + 1)$ and this says that $p \mid (\frac{t_{23}}{t_{22}})^2 + 1$ which can happen if and only if $p \equiv 1 \pmod{4}$, and so we assume that this is the case. By Hensel's lemma there exists $i \in \mathbb{Z}_p$ such that $i^2 + 1 = 0$. Then condition (4') is just $t_{22} \mid (t_{23} - it_{22})(t_{23} + it_{22})$ which is equivalent to $t_{22}t_{33} \mid t_{23} - it_{22}$ or $t_{22}t_{33} \mid t_{23} + it_{22}$. It is easy to see that these cases split our domain in two disjoint subsets. We do the integral over each of these subsets, according to Remark 2.2, and we do the sum of them. The result is

$$\begin{aligned} & 2(1 - p^{-1})^{-3} \int_{|t_{22}| > |t_{33}|} |t_{11}|^{s-1} |t_{22}|^{s-1} |t_{33}|^{s-3} d\mu \\ & = 2p^{1-s} \zeta_p(s) \zeta_p(s - 1) \zeta_p(2s - 2) \end{aligned}$$

Summing the results obtained in the two cases we can obtain the expression of the integral (2.8) for any $p \neq 2$ using χ_4 :

$$\begin{aligned}\zeta_{\mathfrak{G}_4, N_0, p}^{\leq}(s) &= \zeta_p(s) \zeta_p(2s-2) \left(1 + (\chi_4(p) + 1) \frac{p^{1-s}}{1-p^{1-s}} \right) \\ &= \zeta_p(s) \zeta_p(2s-2) \frac{1 + \chi_4(p)p^{1-s}}{1-p^{1-s}} \\ &= \zeta_p(s) \zeta_p(2s-2) \frac{1-p^{2-2s}}{1-p^{1-s}} \frac{1}{1-\chi_4(p)p^{1-s}} \\ &= \zeta_p(s) \zeta_p(s-1) L(s-1, \chi_4, p).\end{aligned}$$

Doing the product of all the local factors we obtain

$$(2.9) \quad \zeta_{\mathfrak{G}_3, N_0}^{\leq}(s) = \zeta(s) \zeta(s-1) L(s-1, \chi_4) \frac{1}{\zeta_2(s) L(s-1, \chi_4, 2)}$$

which has abscissa of convergence 2. The functional equation for almost all primes is:

$$(2.10) \quad \zeta_{\mathfrak{G}_4, N_0, p}^{\leq}(s)|_{p \rightarrow p^{-1}} = (-1)^3 \chi_4(p) p^{-3s+2} \zeta_{\mathfrak{G}_4, N_0, p}^{\leq}(s).$$

2.2.4. *The Bieberbach group $\mathfrak{G}_5$: $IT=(3, 169)$.* This is the group

$$\mathfrak{G}_5 = \langle \gamma, x_1, x_2, x_3 : [x_i, x_j] = 1, \gamma^6 = 1, \gamma x_2 \gamma^{-1} = x_3, \gamma x_3 \gamma^{-1} = x_2^{-1} x_3 \rangle$$

with Fitting subgroup $N_0 = \langle x_1, x_2, x_3 \rangle$. We have

$$(2.11) \quad \zeta_{\mathfrak{G}_5, N_0, p}^{\leq}(s) = (1-p^{-1})^{-3} \int_{\mathcal{T}'} |t_{11}|^{s-2} |t_{22}|^{s-3} |t_{33}|^{s-4} d\mu$$

where $\mathcal{T}'$ is the set of pairs $(\mathbf{t}, \mathbf{v}) \in Tr(3, \mathbb{Z}_p) \times \mathbb{Z}_p^3$ satisfying:

$$\gamma \mathbf{x}^{\mathbf{v}} \mathbf{x}^{\mathbf{t}_i} (\gamma \mathbf{x}^{\mathbf{v}})^{-1} \in B_{\mathbf{t}} \text{ for } i = 1, 2, 3 \text{ and } (\gamma \mathbf{x}^{\mathbf{v}})^6 \in B_{\mathbf{t}}.$$

These conditions are translated in:

$$x_2^{-t_{12}-t_{13}} x_3^{t_{12}} \in B_{\mathbf{t}}, \quad x_2^{-t_{23}} x_3^{t_{22}+t_{23}} \in B_{\mathbf{t}}, \quad x_2^{t_{33}} \in B_{\mathbf{t}} \text{ and } x_1^{6v_1+1} \in B_{\mathbf{t}}.$$

The associated cone conditions are:

- (1) $t_{22} | t_{12} + t_{13};$
- (2) $t_{33} | \frac{t_{12}+t_{13}}{t_{22}} t_{23} + t_{12};$
- (3) $t_{22} | t_{23};$
- (4) $t_{33} | \frac{t_{23}}{t_{22}} t_{23} + t_{22} + t_{23};$
- (5) $t_{22} | t_{33};$
- (6) $t_{33} | \frac{t_{33}}{t_{22}} t_{23};$
- (7) $t_{11} | 6v_1 + 1;$
- (8) $t_{22} | -\frac{6v_1+1}{t_{11}} t_{12};$
- (9) $t_{33} | \frac{6v_1+1}{t_{11}} (\frac{t_{12}}{t_{22}} t_{23} - t_{13}).$

As in the other computations we can see that (3) implies (6), (1) and (8) can be replaced by $t_{22}|t_{12}$ and $t_{22}|t_{13}$, and conditions (4) and (5) imply (3). Similarly, it can be proved that (2) and (9) can be replaced by $t_{33}|\frac{t_{12}}{t_{22}}t_{23} - t_{13}$ and this new condition with (5) imply that $t_{22}|t_{13}$.

The new set of conditions is

- (1') $t_{22}|t_{33}$;
- (2') $t_{22}|t_{12}$;
- (3') $t_{33}|\frac{t_{23}}{t_{22}}t_{23} + t_{22} + t_{23}$;
- (4') $t_{11}|6v_1 + 1$ (or $t_{11} \in \mathbb{Z}_p^*$ for $p = 2, 3$);
- (5') $t_{33}|\frac{t_{12}}{t_{22}}t_{23} - t_{13}$.

If $p = 2$ then it is easy to see that (3') holds if and only if $|t_{22}| = |t_{33}|$ and $t_{22}|t_{23}$. The integral under this new set of conditions is

$$\zeta_{\mathfrak{S}_5, N_0, 2}^{\leq}(s) = (1 - 2^{-1})^{-3} \int_{t_{11} \in \mathbb{Z}_2^*, |t_{22}| = |t_{33}|} |t_{22}|^{s-1} |t_{33}|^{s-3} d\mu = \zeta_2(2s - 2).$$

If $p = 3$ then as in the other cases it is easy to see that (3') holds if and only if $|t_{22}| = |t_{33}|$ and $t_{22}|t_{23}$, or $|t_{22}| = 3|t_{33}|$ and $3t_{22}|t_{23} - t_{22}$. The integral in both cases are respectively

$$(1 - 3^{-1})^{-3} \int_{t_{11} \in \mathbb{Z}_3^*, |t_{22}| = |t_{33}|} |t_{22}|^{s-1} |t_{33}|^{s-3} d\mu = \zeta_3(2s - 2)$$

and

$$(1 - 3^{-1})^{-3} \int_{t_{11} \in \mathbb{Z}_3^*, |t_{22}| = 3|t_{33}|} 3^{-1} |t_{22}|^{s-1} |t_{33}|^{s-3} d\mu = 3^{1-s} \zeta_3(2s - 2).$$

Thus,

$$\zeta_{\mathfrak{S}_5, N_0, 3}^{\leq}(s) = (1 + 3^{1-s}) \zeta_3(2s - 2) = \zeta_2(s - 1).$$

Suppose that $p \neq 2, 3$. We first consider the case $|t_{22}| = |t_{33}|$ where condition (3') can be changed by $t_{22}|t_{23}$. The integral is

$$(1 - p^{-1})^{-3} \int_{|t_{22}| = |t_{33}|} |t_{11}|^{s-1} |t_{22}|^{s-1} |t_{33}|^{s-3} d\mu = \zeta_p(s) \zeta_p(2s - 2)$$

The case $|t_{22}| > |t_{33}|$ and condition (3') imply that $\frac{t_{23}}{t_{22}}$ is a solution of $x^2 + x + 1 \equiv 0 \pmod{p}$ which can happen only if $p \equiv 1 \pmod{3}$ and so we assume that this is the case. By Hensel's lemma there exists $\eta_1, \eta_2 \in \mathbb{Z}_p$ such that $x^2 + x + 1 = (x - \eta_1)(x - \eta_2)$. It follows that condition (3') is equivalent to $t_{22}|(t_{23} - \eta_1 t_{22})(t_{23} - \eta_2 t_{22})$, which is equivalent to the condition: $t_{22}|t_{23} - \eta_1 t_{22}$ or $t_{22}|t_{23} - \eta_2 t_{22}$, and these cases split the

domain in two disjoint subsets. We compute the integral in both cases. The sum of them is

$$2(1 - p^{-1})^{-3} \int_{|t_{22}| > |t_{33}|} |t_{11}|^{s-1} |t_{22}|^{s-1} |t_{33}|^{s-3} d\mu = 2p^{1-s} \zeta_p(s) \zeta_p(s-1) \zeta_p(2s-2).$$

Observe that this result is the same as the one obtained for the local zeta functions of $\mathfrak{G}_3$ for $p \neq 2, 3$ and so we will write just the result:

$$\zeta_{\mathfrak{G}_5, N_0, p}^{\leq}(s) = \zeta_p(s) \zeta_p(s-1) L(s-1, \chi_3, p).$$

Doing the product of all local factors we obtain

$$(2.12) \quad \zeta_{\mathfrak{G}_5, N_0}^{\leq}(s) = \zeta(s) \zeta(s-1) L(s-1, \chi_3).$$

$$(2.13) \quad \frac{\zeta_2(2s-2)}{\zeta_2(s) \zeta_2(s-1) L(s-1, \chi_3, 2)} \frac{1}{\zeta_3(s) L(s-1, \chi_3, 3)}$$

which has abscissa of convergence 2. The functional equation for almost all primes is

$$(2.14) \quad \zeta_{\mathfrak{G}_5, N_0, p}^{\leq}(s)|_{p \rightarrow p^{-1}} = (-1)^3 \chi_3(p) p^{-3s+2} \zeta_{\mathfrak{G}_5, N_0, p}^{\leq}(s).$$

2.2.5. *The Bieberbach group $\mathfrak{G}_6$: $IT=(3,19)$.* This is the group

$$\mathfrak{G}_6 = \langle \alpha, \beta, x_1, x_2, x_3 : [x_i, x_j] = 1, \alpha^2 = x_1, \alpha x_2 \alpha^{-1} = x_2^{-1}, \alpha x_3 \alpha^{-1} = x_3^{-1}, \beta^2 = x_2, \beta x_1 \beta^{-1} = x_1^{-1}, \beta x_3 \beta^{-1} = x_3^{-1}, (\alpha \beta)^2 = x_3^{-1} \rangle$$

with Fitting subgroup $N_0 = \langle x_1, x_2, x_3 \rangle$. We have

$$(2.15) \quad \zeta_{\mathfrak{G}_6, N_0, p}^{\leq}(s) = (1 - p^{-1})^{-3} \int_{\mathcal{T}'} |t_{11}|^{s-3} |t_{22}|^{s-4} |t_{33}|^{s-5} d\mu$$

where $\mathcal{T}'$ is the set of pairs $(\mathbf{t}, \mathbf{v}) \in Tr(3, \mathbb{Z}_p) \times M_{3 \times 3}(\mathbb{Z}_p)$ such that

$$\begin{aligned} \alpha \mathbf{x}^{\mathbf{v}_1} \mathbf{x}^{\mathbf{t}_i} (\alpha \mathbf{x}^{\mathbf{v}_1})^{-1} &\in B_{\mathbf{t}} \quad \beta \mathbf{x}^{\mathbf{v}_2} \mathbf{x}^{\mathbf{t}_i} (\beta \mathbf{x}^{\mathbf{v}_2})^{-1} \in B_{\mathbf{t}} \text{ for } i = 1, 2, 3 \\ (\alpha \mathbf{x}^{\mathbf{v}_1})^2, (\beta \mathbf{x}^{\mathbf{v}_2})^2, (\alpha \mathbf{x}^{\mathbf{v}_1} \beta \mathbf{x}^{\mathbf{v}_2})^2 &\in B_{\mathbf{t}}. \end{aligned}$$

These conditions are translated in

$$x_2^{2t_{12}} x_3^{2t_{13}} \in B_{\mathbf{t}}, \quad x_3^{2t_{13}}, x_3^{2t_{23}} \in B_{\mathbf{t}}, \quad x_1^{2v_{11}+1}, x_2^{2v_{12}}, x_3^{-1-2v_{13}+2v_{23}} \in B_{\mathbf{t}};$$

and the associated cone conditions are:

- (1) $t_{22} | 2t_{12};$
- (2) $t_{33} | \frac{2t_{12}}{t_{22}} t_{23};$
- (3) $t_{33} | 2t_{13};$
- (4) $t_{33} | 2t_{23};$
- (5) $t_{11} | 2v_{11} + 1;$
- (6) $t_{22} | \frac{2v_{11}+1}{t_{11}} t_{12};$
- (7) $t_{33} | \frac{2v_{11}+1}{t_{11}} (\frac{t_{12}}{t_{22}} t_{23} - t_{13});$
- (8) $t_{22} | 2v_{22} + 1;$

- (9) $t_{33} \mid \frac{2v_{22}+1}{t_{22}} t_{23}$;
- (10) $t_{33} \mid -1 - 2v_{13} + 2v_{23}$.

When $p \neq 2$ these conditions are clearly equivalent to

$$t_{22} \mid t_{12}, \quad t_{33} \mid t_{13}, \quad t_{33} \mid t_{23}, \quad t_{11} \mid 2v_{11} + 1, \quad t_{22} \mid 2v_{22} + 1, \quad t_{33} \mid -1 - 2v_{13} + 2v_{23};$$

and by Remark 2.2 the integral under this set of conditions is

$$\zeta_{\mathfrak{B}_1, N_0, p}^{\leq}(s) = (1 - p^{-1})^{-3} \int_{\mathcal{T}'} |t_{11}|^{s-2} |t_{22}|^{s-2} |t_{33}|^{s-2} d\mu = \zeta_p(s-1)^3$$

When $p = 2$, then conditions (5), (8) y (10) imply that t_{11}, t_{22} and t_{33} are units and therefore all conditions are trivially satisfied. Then the integral (2.15) for $p = 2$ is equal to 1. Doing the product of all the local factors we obtain

$$(2.16) \quad \zeta_{\mathfrak{G}_6, N_0}^{\leq}(s) = \zeta(s-1)^3 \zeta_2(s-1)^{-3},$$

which has abscissa of convergence 2. The functional equation for almost all primes is

$$(2.17) \quad \zeta_{\mathfrak{G}_6, N_0, p}^{\leq}(s)|_{p \rightarrow p^{-1}} = (-1)^3 p^{-3s+3} \zeta_{\mathfrak{G}_6, N_0, p}^{\leq}(s).$$

2.2.6. *The Bieberbach group $\mathfrak{B}_1$: $IT=(3,7)$.* This is the group

$$\mathfrak{B}_1 = \langle \gamma, x_1, x_2, x_3 : [x_i, x_j] = 1, \gamma^2 = x_1, \gamma x_2 \gamma^{-1} = x_2, \gamma x_3 \gamma^{-1} = x_3^{-1} \rangle$$

with Fitting subgroup $N_0 = \langle x_1, x_2, x_3 \rangle$. We have

$$(2.18) \quad \zeta_{\mathfrak{B}_1, N_0, p}^{\leq}(s) = (1 - p^{-1})^{-3} \int_{\mathcal{T}'} |t_{11}|^{s-2} |t_{22}|^{s-3} |t_{33}|^{s-4} d\mu$$

where $\mathcal{T}'$ is the set of pairs $(\mathbf{t}, \mathbf{v}) \in Tr(3, \mathbb{Z}_p) \times \mathbb{Z}_p^3$ satisfying the conditions:

$$(\gamma \mathbf{x}^{\mathbf{v}}) \mathbf{x}^{\mathbf{t}_i} (\gamma \mathbf{x}^{\mathbf{v}})^{-1} \in B_{\mathbf{t}} \text{ for } i = 1, 2, 3 \text{ and } (\gamma \mathbf{x}^{\mathbf{v}})^2 \in B_{\mathbf{t}}.$$

These conditions are equivalent to

$$x_3^{2t_{13}}, x_3^{2t_{23}} \in B_{\mathbf{t}} \text{ and } x_1^{2v_1+1} x_2^{2v_2} \in B_{\mathbf{t}}.$$

The associated cone conditions are:

- (1) $t_{33} \mid 2t_{13}$;
- (2) $t_{33} \mid 2t_{23}$;
- (3) $t_{11} \mid 2v_1 + 1$;
- (4) $t_{22} \mid -\frac{2v_1+1}{t_{11}} t_{12} + 2v_2$;
- (5) $t_{33} \mid -\frac{-\frac{2v_1+1}{t_{11}} t_{12} + 2v_2}{t_{22}} t_{23} - \frac{2v_1+1}{t_{11}} t_{13}$.

When $p \neq 2$ condition (1) and (2) are just $t_{33}|t_{13}$ and $t_{33}|t_{23}$ and they imply condition (5). Then the conditions are

$$t_{33}|t_{13}, t_{33}|t_{23}, t_{11}|2v_1 + 1 \text{ and } t_{22}| -\frac{2v_1 + 1}{t_{11}}t_{12} + 2v_2.$$

Therefore, by Remark 2.2, the integral (2.18) is:

$$\zeta_{\mathfrak{B}_1, N_0, p}^{\leq}(s) = (1 - p^{-1})^{-3} \int |t_{11}|^{s-1} |t_{22}|^{s-2} |t_{33}|^{s-2} d\mu = \zeta_p(s) \zeta_p(s-1)^2.$$

When $p = 2$ then condition (3) says that t_{11} is a unit. We shall consider cases:

Case 1.1: $t_{33}|t_{13}$ and $t_{33}|t_{23}$. In this case condition (5) is redundant and therefore the conditions are: $t_{11} \in \mathbb{Z}_2^*$, $t_{33}|t_{13}$, $t_{33}|t_{23}$ and $t_{22}|t_{12} - \frac{2v_2 t_{11}}{2v_1 + 1}$ and according to Remark 2.2 the integral under these conditions is

$$(1 - 2^{-1})^{-3} \int_{t_{11} \in \mathbb{Z}_2^*} |t_{22}|^{s-2} |t_{33}|^{s-2} d\mu = \zeta_2(s-1)^2.$$

Case 1.2: $t_{33}|t_{13}$ and $|t_{33}| = |2t_{23}|$. Observe that necessarily $t_{33} \in 2\mathbb{Z}_2$. The coefficient of t_{23} in (5) must lie in $2\mathbb{Z}_2$ because otherwise (5) and $t_{33}|t_{13}$ would imply $t_{33}|t_{23}$. Then our conditions are: $t_{11} \in \mathbb{Z}_2^*$, $t_{33} \in 2\mathbb{Z}_2$, $t_{33}|t_{13}$, $|2^{-1}t_{33}| = |t_{23}|$, $2t_{22}|t_{12} - \frac{2v_2 t_{11}}{2v_1 + 1}$ and the integral under these conditions is

$$2^{-1}(1 - 2^{-1})^{-3} \int_{t_{11} \in \mathbb{Z}_2^*, t_{33} \in 2\mathbb{Z}_2} 2^{-1} |t_{22}|^{s-2} 2 |t_{33}|^{s-2} d\mu = 2^{-s} \zeta_2(s-1)^2.$$

Case 2.1: $|t_{33}| = |2t_{13}|$ y $t_{33}|t_{23}$. By condition (5) this would imply that $t_{33}|t_{13}$ which is a contradiction. Then this case is impossible.

Case 2.2: $|t_{33}| = |2t_{13}| = |2t_{23}|$. Since the coefficient of t_{13} in (5) is a unit then this also implies that the coefficient of t_{23} is a unit and therefore we have $|t_{22}| = |t_{12} - \frac{2v_2 t_{11}}{2v_1 + 1}|$ which implies (4). But now condition conditions (1) and (5) reduce (5) into $t_{33}|t_{23} + t_{13}$ and this is implied by the hypothesis because $|t_{13}| = |t_{23}|$ implies that $|t_{13} + t_{23}| < |t_{13}|$ and hence $t_{33}|t_{13} + t_{23}$. Then the conditions are: $t_{11} \in \mathbb{Z}_2^*$, $t_{33} \in 2\mathbb{Z}_2$, $|2^{-1}t_{33}| = |t_{13}|$, $|2^{-1}t_{33}| = |t_{23}|$ and $|t_{22}| = |t_{12} - \frac{2v_2 t_{11}}{2v_1 + 1}|$. The integral under these conditions is

$$2^{-3}(1 - 2^{-1})^{-3} \int_{t_{11} \in \mathbb{Z}_2^*, t_{33} \in 2\mathbb{Z}_2} |t_{22}|^{s-2} 2 |t_{33}|^{s-2} d\mu = 2^{-s} \zeta_2(s-1)^2.$$

Summing these results we obtain:

$$\zeta_{\mathfrak{B}_1, N_0, 2}^{\leq}(s) = (1 + 2^{1-s}) \zeta_2(s-1)^2.$$

Doing the product of all the local factors we obtain

$$(2.19) \quad \zeta_{\mathfrak{B}_1, N_0}^{\leq}(s) = \zeta(s-1)^2 \zeta(s) \zeta_2(s)^{-1} (1 + 2^{1-s}),$$

which has abscissa of convergence 2. The functional equation for almost all primes is:

$$(2.20) \quad \zeta_{\mathfrak{B}_1, N_0, p}^{\leq}(s)|_{p \rightarrow p^{-1}} = (-1)^3 p^{-3s+2} \zeta_{\mathfrak{B}_1, N_0, p}^{\leq}(s).$$

2.2.7. *The Bieberbach group $\mathfrak{B}_2$:* $IT=(3,9)$. This is the group

$$\mathfrak{B}_2 = \langle \gamma, x_1, x_3, x_3 : [x_i, x_j] = 1, \gamma^2 = x_1, \gamma x_2 \gamma^{-1} = x_2, \gamma x_3 \gamma^{-1} = x_2 x_3^{-1} \rangle$$

with Fitting subgroup $N_0 = \langle x_1, x_2, x_3 \rangle$. We have

$$(2.21) \quad \zeta_{\mathfrak{B}_2, N_0, p}^{\leq}(s) = (1 - p^{-1})^{-1} \int_{\mathcal{T}'} |t_{11}|^{s-2} |t_{22}|^{s-3} |t_{33}|^{s-4} d\mu$$

where $\mathcal{T}'$ is the set of pairs $(\mathbf{t}, \mathbf{v}) \in Tr(3, \mathbb{Z}_p) \times \mathbb{Z}_p^3$ such that

$$\gamma \mathbf{x}^{\mathbf{v}} \mathbf{x}^{\mathbf{t}_i} (\gamma \mathbf{x}^{\mathbf{v}})^{-1} \in B_{\mathbf{t}} \text{ for } i = 1, 2, 3 \text{ and } (\gamma \mathbf{x}^{\mathbf{v}})^2 \in B_{\mathbf{t}}.$$

These conditions are translated in:

$$x_2^{t_{13}} x_3^{-2t_{13}}, x_2^{t_{23}} x_3^{-2t_{23}}, x_2^{t_{33}} x_3^{-2t_{33}}, x_1^{2v_1+1} x_2^{2v_2+v_3} \in B_{\mathbf{t}}.$$

The associated cone conditions are:

- (1) $t_{22}|t_{13}$;
- (2) $t_{33}|t_{22}t_{23} - 2t_{13}$;
- (3) $t_{22}|t_{23}$;
- (4) $t_{33}|t_{22}t_{23} - 2t_{23}$;
- (5) $t_{22}|t_{33}$;
- (6) $t_{33}|t_{22}t_{23} - 2t_{33}$;
- (7) $t_{11}|2v_1 + 1$;
- (8) $t_{22}|t_{11} - \frac{2v_1+1}{t_{11}}t_{12} + 2v_2 + v_3$;
- (9) $t_{33}|t_{11} - \frac{2v_1+1}{t_{11}}t_{12} + 2v_2 + v_3$.

Assume that $p \neq 2$. We shall consider cases:

Case 1: $|t_{33}| \geq |t_{13}|, |t_{33}| \geq |t_{23}|$. In this case all conditions are implied by (5), (7) and (8). Then the conditions are just $t_{22}|t_{33}, t_{33}|t_{13}, t_{33}|t_{23}, t_{11}|2v_1 + 1$ and $t_{22}|t_{11} - \frac{2v_1+1}{t_{11}}t_{12} + 2v_2 + v_3$. The integral is

$$(1 - p^{-1})^{-3} \int_{t_{22}|t_{33}} |t_{11}|^{s-1} |t_{22}|^{s-2} |t_{33}|^{s-2} d\mu = \zeta_p(s) \zeta_p(s-1) \zeta_p(2s-2).$$

Case 2: $|t_{13}| > |t_{33}| \geq |t_{23}|$. This condition is impossible because the hypothesis and (2) imply $t_{33}|t_{13}$.

Case 3: $|t_{23}| > |t_{33}| \geq |t_{13}|$. Observe that (3) and (4) imply (1)-(6) and even more (4) implies (3) because $t_{23}|t_{33}$ with (4) imply $1|\frac{t_{23}}{t_{22}}$. Observe also that (8) and (9) can be replaced by (*) $t_{22}|\frac{t_{33}}{t_{23}}| - \frac{2v_1+1}{t_{11}}t_{12} + 2v_2 + v_3$. We shall do a change of variables $t_{33} = pt_{23}a$ and $t_{13} = pt_{23}ab$ and then condition (4) is translated into $pa| - \frac{t_{23}}{t_{22}} - 2$, or $pat_{22}| -$

$t_{23} - 2t_{22}$ and condition (*) is just $pat_{22}| - \frac{2v_1+1}{t_{11}}t_{12} + 2v_2 + v_3$. Taking account of the change of variables we have to compute the integral of $(1 - p^{-1})^{-3}p^{2-s}|t_{11}|^{s-2}|t_{22}|^{s-3}|t_{23}|^{s-2}|a|^{s-3}$ under the following set of conditions:

- (1') $pat_{22}| - t_{23} - 2t_{22}$;
- (2') $t_{11}|2v_1 + 1$ (or $t_{11} \in \mathbb{Z}_2^*$ if $p = 2$);
- (3') $pat_{22}| - \frac{2v_1+1}{t_{11}}t_{12} + 2v_2 + v_3$.

Since $p \neq 2$ then condition (1') implies that in the domain of integration we have $|t_{22}| = |t_{23}|$ and therefore we have just to compute the integral of $(1 - p^{-1})^{-3}p^{2-s}|t_{11}|^{s-2}|t_{22}|^{2s-5}|a|^{s-3}$ under the same domain of integration. The result, according to Remark 2.2, is

$$(1 - p^{-1})^{-3}p^{2-s} \int |t_{11}|^{s-1}|t_{22}|^{2s-3}|a|^{s-1}p^{-2}d\mu = p^{-s}\zeta_p(s)^2\zeta_p(2s-2).$$

Case 4: $|t_{23}| \geq |t_{13}| > |t_{33}|$. As in the last case, condition (4) implies (1)-(6) and since $t_{23}|t_{13}$ and $t_{23}|t_{33}$ then condition (9) implies (8). Doing the change of variables $t_{13} = at_{23}$, $t_{33} = pabt_{23}$ we translate conditions (4) and (9) in $pabt_{22}| - t_{23} - 2t_{22}$ and $t_{22}pab| - (-\frac{2v_1+1}{t_{11}}t_{12} + 2v_2 + v_3) - \frac{2v_1+1}{t_{11}}at_{22}$. Taking account on the change of variables we have to integrate $(1 - p^{-1})^{-3}p^{3-s}|t_{11}|^{s-2}|t_{22}|^{s-3}|a|^{s-3}|b|^{s-4}|t_{23}|^{s-2}$ under the following conditions:

- (1') $pabt_{22}| - t_{23} - 2t_{22}$;
- (2') $t_{11}|2v_1 + 1$ (or $t_{11} \in \mathbb{Z}_2^*$ if $p = 2$);
- (3') $t_{22}pab|\frac{2v_1+1}{t_{11}}t_{12} - 2v_2 - v_3 - \frac{2v_1+1}{t_{11}}at_{22}$.

Since $p \neq 2$ then condition (1') implies $|t_{23}| = |t_{22}|$ and therefore we can replace the integrand by $(1 - p^{-1})^{-3}p^{3-s}|t_{11}|^{s-2}|t_{22}|^{2s-5}|a|^{s-3}|b|^{s-4}$ and compute the integral according to Remark 2.2. The result is

$$\begin{aligned} (1 - p^{-1})^{-3}p^{3-s} \int p^{-2}|t_{11}|^{s-1}|t_{22}|^{2s-3}|a|^{s-1}|b|^{s-2}d\mu = \\ = p^{1-s}\zeta_p(s)^2\zeta_p(s-1)\zeta_p(2s-2)(1 - p^{-1}) \end{aligned}$$

Case 5: $|t_{13}| > |t_{23}| > |t_{33}|$. In this case, as in the last two cases, condition (2) implies (1)-(6) and conditions (8) and (9) imply (7). Doing the change of variables $t_{23} = pat_{13}$, $t_{33} = p^2abt_{13}$ condition (2) becomes into $p^2ab| - pa\frac{t_{23}}{t_{22}} - 2$ which leads a contradiction since $p \neq 2$.

Adding the results we obtain for $p \neq 2$:

$$\zeta_{\mathfrak{B}_2, N_0, p}^{\leq}(s) = \zeta_p(s)\zeta_p(s-1)^2.$$

For $p = 2$ it is not difficult to see that $\mathfrak{B}_2$ has exactly three subgroups of index 2. One of them is N and the others two, say A_1 and A_2 , are isomorphic to $\mathfrak{B}_1$ and they have intersection contained in N . Since

the intersection of subgroups of $\mathfrak{B}_2$ of index a power of 2 has index a power of 2 then every subgroup A of $\mathfrak{B}_2$ of index a power of 2 satisfying $AN = \mathfrak{B}_2$ satisfies one and only one of the following two options: $A \leq A_1$ or $A \leq A_2$. Then

$$\begin{aligned}\zeta_{\mathfrak{B}_2, N_0, 2}^{\leq}(s) &= 1 + 2\zeta_{\mathfrak{B}_2, N, 2}(s) \\ &= \zeta(s-1)^2(1 + 2^{1-s} + 2^{3-2s})\end{aligned}$$

Doing the product of all the local factors we obtain:

$$(2.22) \quad \zeta_{\mathfrak{B}_2, N_0}^{\leq}(s) = \zeta(s)\zeta(s-1)^2 \frac{1 + 2^{1-s} + 2^{3-2s}}{\zeta_2(s)}$$

which has abscissa of convergence 2. The functional equation for almost all primes is:

$$(2.23) \quad \zeta_{\mathfrak{B}_2, N_0, p}^{\leq}(s)|_{p \rightarrow p^{-1}} = (-1)^3 p^{-3s+2} \zeta_{\mathfrak{B}_2, N_0, p}^{\leq}(s).$$

2.2.8. *The Bieberbach group $\mathfrak{B}_3$: $IT=(3, 29)$.* This is the group

$$\begin{aligned}\mathfrak{B}_3 = \langle \alpha, \beta, x_1, x_2, x_3 : [x_i, x_j] = 1, \alpha^2 = x_1, \alpha x_2 \alpha^{-1} = x_2^{-1}, \alpha x_3 \alpha^{-1} = x_3^{-1} \\ \beta^2 = x_2, \beta x_1 \beta^{-1} = x_1, \beta x_3 \beta^{-1} = x_3^{-1}, \beta \alpha \beta^{-1} \alpha^{-1} = x_2 \rangle\end{aligned}$$

with Fitting subgroup $N_0 = \langle x_1, x_2, x_3 \rangle$. We have

$$(2.24) \quad \zeta_{\mathfrak{B}_3, N_0, p}^{\leq}(s) = (1 - p^{-1})^{-1} \int_{\mathcal{T}'} |t_{11}|^{s-3} |t_{22}|^{s-4} |t_{33}|^{s-5} d\mu$$

where $\mathcal{T}'$ is the set of pairs $(\mathbf{t}, \mathbf{v}) \in Tr(3, \mathbb{Z}_p) \times M_{2 \times 3}(\mathbb{Z}_p)$ such that

$$\begin{aligned}\alpha \mathbf{x}^{\mathbf{v}_1} \mathbf{x}^{\mathbf{t}_i} (\alpha \mathbf{x}^{\mathbf{v}_1})^{-1}, \beta \mathbf{x}^{\mathbf{v}_2} \mathbf{x}^{\mathbf{t}_i} (\beta \mathbf{x}^{\mathbf{v}_2})^{-1} &\in B_{\mathbf{t}} \text{ for } i = 1, 2, 3 \\ (\alpha \mathbf{x}^{\mathbf{v}_1})^2, (\beta \mathbf{x}^{\mathbf{v}_2})^2, \beta \mathbf{x}^{\mathbf{v}_2} \alpha \mathbf{x}^{\mathbf{v}_1} (\beta \mathbf{x}^{\mathbf{v}_2})^{-1} &(\alpha \mathbf{x}^{\mathbf{v}_1})^{-1} \in B_{\mathbf{t}}.\end{aligned}$$

These conditions are translated in:

$$x_2^{t_{12}}, x_2^{t_{22}}, x_3^{t_{13}}, x_3^{t_{23}} \in B_{\mathbf{t}}; x_1^{2v_{11}+1}, x_1^{2v_{21}} x_2^{2v_{22}+1}, x_2^{2v_{22}+1} x_3^{2v_{13}-2v_{23}} \in B_{\mathbf{t}}.$$

The associated cone conditions are:

- (1) $t_{22} | 2t_{12};$
- (2) $t_{33} | \frac{2t_{12}}{t_{22}} t_{23};$
- (3) $t_{33} | 2t_{13};$
- (4) $t_{33} | 2t_{23};$
- (5) $t_{11} | 2v_{11} + 1;$
- (6) $t_{22} | -\frac{2v_{11}+1}{t_{11}};$
- (7) $t_{33} | \frac{2v_{11}+1}{t_{11}} (\frac{t_{12}}{t_{22}} t_{23} - t_{13});$
- (8) $t_{11} | 2v_{21};$
- (9) $t_{22} | -\frac{2v_{21}}{t_{11}} t_{12} + 2v_{22} + 1;$
- (10) $t_{33} | -\frac{-\frac{2v_{21}}{t_{11}} t_{12} + 2v_{22} + 1}{t_{22}} t_{23} - \frac{2v_{21}}{t_{11}} t_{13};$

- (11) $t_{22}|2v_{22} + 1;$
 (12) $t_{33}| -\frac{2v_{22}+1}{t_{22}}t_{23} + 2v_{13} - 2v_{23}.$

When $p \neq 2$ then it is easy to see that these conditions are reduced to:

$$t_{22}|t_{12}, t_{33}|t_{13}, t_{33}|t_{23}, t_{11}|2v_{11} + 1, t_{11}|2v_{21}, t_{22}|2v_{22} + 1 \text{ and } t_{33}|v_{13} - v_{23}.$$

The integral under these conditions is:

$$\zeta_{\mathfrak{B}_3, N_0, p}^{\leq}(s) = (1 - p^{-1})^{-3} \int |t_{11}|^{s-1} |t_{22}|^{s-2} |t_{33}|^{s-2} d\mu = \zeta_p(s) \zeta_p(s-1)^2.$$

When $p = 2$, (5) and (11) say that t_{11} and t_{22} are units. Combining this fact with (3) and (4) we can see from (10) that $t_{33}|t_{23}$, and using (7) we obtain $t_{33}|t_{13}$. The reduced conditions are therefore:

$$t_{11} \in \mathbb{Z}_2^*, t_{22} \in \mathbb{Z}_2^*, t_{33}|t_{13}, t_{33}|t_{23} \text{ and } t_{33}|2v_{13} - 2v_{23}.$$

We split the domain of integration according to when $t_{33} \in \mathbb{Z}_2^*$ or $t_{33} \in 2\mathbb{Z}_2$. In the later case the condition $t_{33}|2v_{13} - 2v_{23}$ is equivalent to $\frac{t_{33}}{2}|v_{13} - v_{23}$. The results are respectively:

$$(1 - 2^{-1})^{-3} \int_{t_{11}, t_{22}, t_{33} \in \mathbb{Z}_2^*} 1 d\mu = 1$$

and

$$(1 - 2^{-1})^{-3} \int_{t_{11}, t_{22} \in \mathbb{Z}_2^*, t_{33} | 2\mathbb{Z}_2} 2|t_{33}|^{s-2} d\mu = 2^{2-s} \zeta_2(s-1)$$

Therefore

$$\zeta_{\mathfrak{B}_3, N_0, 2}^{\leq}(s) = 1 + 2^{2-s} \zeta_2(s-1) = (1 + 2^{1-s}) \zeta_2(s-1).$$

The product of all the local factors is

$$(2.25) \quad \zeta_{\mathfrak{B}_3, N_0}^{\leq}(s) = \zeta(s) \zeta(s-1)^2 \frac{1 + 2^{1-s}}{\zeta_2(s) \zeta_2(s-1)},$$

which has abscissa of convergence 2. The functional equation for almost all primes is:

$$(2.26) \quad \zeta_{\mathfrak{B}_3, N_0, p}^{\leq}(s)|_{p \rightarrow p^{-1}} = (-1)^3 p^{-3s+2} \zeta_{\mathfrak{B}_3, N_0, p}^{\leq}(s).$$

2.2.9. *The Bieberbach group $\mathfrak{B}_4$: $IT(3,33)$.* This is the group

$$\mathfrak{B}_4 = \langle \alpha, \beta, x_1, x_2, x_3 : [x_i, x_j] = 1, \alpha^2 = x_1, \alpha x_2 \alpha^{-1} = x_2^{-1}, \alpha x_3 \alpha^{-1} = x_3^{-1}, \beta^2 = x_2, \beta x_1 \beta^{-1} = x_1, \beta x_3 \beta^{-1} = x_3^{-1}, \beta \alpha \beta^{-1} \alpha^{-1} = x_2 x_3 \rangle$$

with Fitting subgroup $N_0 = \langle x_1, x_2, x_3 \rangle$. We have

$$(2.27) \quad \zeta_{\mathfrak{B}_4, N_0, p}^{\leq}(s) = (1 - p^{-1})^{-3} \int_{\mathcal{T}'} |t_{11}|^{s-3} |t_{22}|^{s-4} |t_{33}|^{s-5} d\mu$$

where $\mathcal{T}'$ is the set of pairs $(\mathbf{t}, \mathbf{v}) \in Tr(3, \mathbb{Z}_p) \times M_{2 \times 3}(\mathbb{Z}_p)$ satisfying:

$$\begin{aligned} \alpha \mathbf{x}^{\mathbf{v}_1} \mathbf{x}^{\mathbf{t}_i} (\alpha \mathbf{x}^{\mathbf{v}_1})^{-1}, \beta \mathbf{x}^{\mathbf{v}_2} \mathbf{x}^{\mathbf{t}_i} (\alpha \mathbf{x}^{\mathbf{v}_2})^{-1} &\in B_{\mathbf{t}} \text{ for } i = 1, 2, 3 \\ (\alpha \mathbf{x}^{\mathbf{v}_1})^2, (\beta \mathbf{x}^{\mathbf{v}_2})^2, \beta \mathbf{x}^{\mathbf{v}_1} \alpha \mathbf{x}^{\mathbf{v}_2} (\beta \mathbf{x}^{\mathbf{v}_1})^{-1} (\alpha \mathbf{x}^{\mathbf{v}_2})^{-1} &\in B_{\mathbf{t}}. \end{aligned}$$

These conditions are translated in:

$$x_2^{2t_{i2}}, x_3^{2t_{i3}} \in B_{\mathbf{t}} \text{ for } i = 1, 2, 3, \quad x_1^{2v_{11}+1}, x_1^{2v_{21}} x_2^{2v_{22}+1} \in B_{\mathbf{t}}.$$

The associated cone conditions are:

- (1) $t_{22} | 2t_{12};$
- (2) $t_{33} | \frac{2t_{12}}{t_{22}} t_{23};$
- (3) $t_{33} | 2t_{23};$
- (4) $t_{33} | 2t_{13};$
- (5) $t_{11} | 2v_{11} + 1;$
- (6) $t_{22} | \frac{2v_{11}+1}{t_{11}} t_{12};$
- (7) $t_{33} | \frac{2v_{11}+1}{t_{11}} (\frac{t_{12}}{t_{22}} t_{23} - t_{13});$
- (8) $t_{11} | 2v_{21};$
- (9) $t_{22} | -\frac{2v_{21}}{t_{11}} t_{12} + 2v_{22} + 1;$
- (10) $t_{33} | -\frac{-\frac{2v_{21}}{t_{11}} t_{12} + 2v_{22} + 1}{t_{22}} t_{23} - \frac{2v_{21}}{t_{11}} t_{13};$
- (11) $t_{22} | 2v_{22} + 1$
- (12) $t_{33} | -\frac{2v_{22}+1}{t_{22}} t_{23} + 2v_{13} - 2v_{23} + 1.$

When $p \neq 2$ then conditions (1), (3), (4) and (8) are just $t_{22} | t_{12}$, $t_{33} | t_{23}$, $t_{33} | t_{13}$ and $t_{11} | v_{21}$. These conditions imply conditions (2), (6), (7) and (10) and simplify conditions (9) and (12). The new set of conditions is

$$t_{11} | v_{21}, t_{22} | t_{12}, t_{33} | t_{13}, t_{33} | t_{23} t_{11} | 2v_{11} + 1, t_{33} | 2v_{13} - 2v_{23} + 1;$$

and the integral (2.27) becomes into

$$\zeta_{\mathfrak{B}_4, N_0, p}^{\leq}(s) = (1 - p^{-1})^{-3} \int |t_{11}|^{s-1} |t_{22}|^{s-2} |t_{33}|^{s-2} d\mu = \zeta_p(s) \zeta_p(s-1)^2.$$

When $p = 2$ then condition (5) says that t_{11} is a unit, and this in combination with (9) imply that t_{22} is a unit. Now, (4), (10) and the fact that t_{11} and t_{22} are units imply that $t_{33} | t_{23}$ and combining this with (12) we obtain that t_{33} is a unit. Then

$$\zeta_{\mathfrak{B}_4, N_0, 2}^{\leq}(s) = (1 - p^{-1})^{-3} \int_{t_{11}, t_{22}, t_{33} \in \mathbb{Z}_2^*} d\mu = 1.$$

Therefore the product of all the local factors is

$$(2.28) \quad \zeta_{\mathfrak{B}_4, N_0}^{\leq}(s) = \zeta(s) \zeta(s-1)^2 \zeta_2(s)^{-1} \zeta_2(s-1)^{-2},$$

which has abscissa of convergence 2. The functional equation for almost all primes is

$$(2.29) \quad \zeta_{\mathfrak{B}_4, N_0, p}^{\leq}(s)|_{p \rightarrow p^{-1}} = (-1)^3 p^{-3s+2} \zeta_{\mathfrak{B}_4, N_0, p}^{\leq}(s)$$

Now we start with the AB -groups which are note Bieberbach groups.

2.2.10. *The AB -groups of the type $Q = p2$.* These are the groups

$$E : \langle x_1, x_2, x_3, \gamma : [x_2, x_1] = x_3^{2q}, [x_3, x_1] = [x_3, x_2] = [x_3, \gamma] = 1, \\ \gamma x_1 = x_1^{-1} \gamma, \gamma x_2 = x_2^{-1} \gamma, \gamma^2 = x_3 \rangle, \quad q \in \mathbb{N}$$

The Fitting subgroup of E is $N = N_{2q} = \langle x_1, x_2, x_3 \rangle$ and we have:

$$(2.30) \quad \zeta_{E, N, p}^{\leq}(s) = (1 - p^{-1})^{-3} \int_{\mathcal{T}'} |t_{11}|^{s-2} |t_{22}|^{s-3} |t_{33}|^{s-4} s \mu$$

where $\mathcal{T}'$ is the set of pairs $(\mathbf{t}, \mathbf{v}) \in Tr(3, \mathbb{Z}_p) \times M_{1 \times 3}(\mathbb{Z}_p)$ satisfying:

$$t_{33} | 2qt_{11}t_{22}, (\gamma \mathbf{x}^{\mathbf{v}})^{-1} \mathbf{x}^{\mathbf{t}_i} \gamma \mathbf{x}^{\mathbf{v}} \in B_{\mathbf{t}} \text{ for } i = 1, 2, 3, \text{ and } \gamma \mathbf{x}^{\mathbf{v}} \gamma \mathbf{x}^{\mathbf{v}} \in B_{\mathbf{t}}.$$

These conditions are translated in:

$$t_{33} | 2qt_{11}t_{22}, x_3^{2t_{13}+2q(t_{11}v_2-v_1t_{12})-2qt_{11}t_{12}} \in B_{\mathbf{t}}, x_3^{2t_{23}+2q(-v_1t_{22})} \in B_{\mathbf{t}} \\ x_3^{2v_3-2qv_1v_2+1} \in B_{\mathbf{t}}.$$

The associated cone conditions are:

- (1) $t_{33} | 2kt_{11}t_{22}$
- (2) $t_{33} | 2t_{13} + 2k(t_{11}v_2 - v_1t_{12}) - 2kt_{11}t_{12}$
- (3) $t_{33} | 2t_{23} - 2kv_1t_{22}$
- (4) $t_{33} | 2v_3 - 2kv_1v_2 + 1$

When $p \neq 2$ then the integral (2.30) can be computed directly using Remark 2.2 and the result is

$$\zeta_{E, N, p}^{\leq}(s) = (1 - p^{-1})^{-3} \int_{t_{33} | kt_{11}t_{22}} |t_{11}|^{s-2} |t_{22}|^{s-3} |t_{33}|^{s-1} d\mu \\ = \zeta_p(s) (\zeta_p(s-1) \zeta_p(s-2) - p^{-s(v_p(k)+1)} \zeta_p(2s-1) \zeta_p(2s-2)).$$

If $p = 2$, then (4) says that t_{33} is a unit and thus the other conditions are trivial. Then integral (2.30) is:

$$\zeta_{E, N, 2}^{\leq}(s) = (1 - 2^{-1})^{-3} \int_{t_{33} \in \mathbb{Z}_2^*} |t_{11}|^{s-2} |t_{22}|^{s-3} d\mu = \zeta_2(s-1) \zeta_2(s-2)$$

Doing the product of all the local factors we obtain:

$$\begin{aligned} \zeta_{E,N}^{\leq}(s) &= \frac{\zeta_2(3s-3)}{\zeta_2(2s-1)\zeta_2(2s-2)} \cdot \\ &\cdot \prod_{p \neq 2, p|q} \frac{\zeta_p(s-1)\zeta_p(s-2) - p^{-s(v_p(q)+1)}\zeta_p(2s-1)\zeta_p(2s-2)}{\zeta_p(s-1)\zeta_p(s-2) - p^{-s}\zeta_p(2s-1)\zeta_p(2s-2)} \\ &\cdot \frac{\zeta(s-1)\zeta(s-2)\zeta(2s-1)\zeta(2s-2)}{\zeta(3s-3)}. \end{aligned}$$

which has abscissa of convergence 3. The functional equation for these groups for almost all primes is:

$$(2.31) \quad \zeta_{E,N,p}^{\leq}(s)|_{p \rightarrow p^{-1}} = (-1)^3 p^{-3s+3} \zeta_{E,N,p}^{\leq}(s).$$

2.3. The AB-groups of type $Q = pg$. These are the groups

$$\begin{aligned} E : \langle x_1, x_2, x_3, \gamma : [x_2, x_1] = x_3^{2q}, [x_3, x_1] = [x_3, x_2] = [\gamma, x_1] = 1, \\ \gamma x_3 = x_3^{-1}\gamma, \gamma x_2 = x_2^{-1}\gamma x_3^{-q}, \gamma^2 = x_1 \rangle, \quad q \in \mathbb{N} \end{aligned}$$

The Fitting subgroup of E is $N = N_{2q} = \langle x_1, x_2, x_3 \rangle$. We have

$$(2.32) \quad \zeta_{E_{2q}, N_{2q}, p}^{\leq}(s) = (1 - p^{-1})^{-3} \int_{\mathcal{T}'} |t_{11}|^{s-2} |t_{22}|^{s-3} |t_{33}|^{s-4} s \mu$$

where $\mathcal{T}'$ is the set of pairs $(\mathbf{t}, \mathbf{v}) \in Tr(3, \mathbb{Z}_p) \times M_{1 \times 3}(\mathbb{Z}_p)$ such that

$$t_{33}|2qt_{11}t_{22}, \quad (\gamma \mathbf{x}^{\mathbf{v}})^{-1} \mathbf{x}^{\mathbf{t}_i} \gamma \mathbf{x}^{\mathbf{v}} \in B_{\mathbf{t}}, \text{ for } i = 1, 2, 3 \text{ and } (\gamma \mathbf{x}^{\mathbf{v}})^2 \in B_{\mathbf{t}}.$$

These conditions are translated in:

$$\begin{aligned} t_{33}|2qt_{11}t_{22}, \quad x_2^{2t_{12}} x_3^{2t_{13}+q+2q(t_{11}v_2+v_1t_{12})}, x_3^{q+2q(v_1t_{22})} \in B_{\mathbf{t}}, \text{ for } i = 1, 2, 3 \\ \text{and } x_1^{2v_1+1} x_3^{-k-2kv_1v_2} \in B_{\mathbf{t}}, \end{aligned}$$

whose associated cone conditions are:

- (1) $t_{33}|2qt_{11}t_{22}$
- (2) $t_{22}|2t_{12}$
- (3) $t_{33}| -\frac{2t_{12}}{t_{22}}t_{23} + 2t_{13} + qt_{12} + 2q(t_{11}v_2 + v_1t_{12})$
- (4) $t_{33}|qt_{22} + 2qv_1t_{22}$
- (5) $t_{11}|2v_1 + 1$
- (6) $t_{22}|\frac{2v_1+1}{t_{11}}t_{12}$
- (7) $t_{33}| -\frac{(2v_1+1)t_{12}}{t_{11}t_{22}}t_{23} + qv_2 + 2qv_1v_2 + \frac{2v_1+1}{t_{11}}t_{13} + q\frac{2v_1+1}{t_{11}}(\frac{2v_1+1}{t_{11}} - 1)t_{11}t_{12}$

If $p \neq 2$ then (2) implies (6), (2) and (5) imply (4), and using (2) and (5) we can reduce (3) in $t_{33}| -\frac{2t_{12}}{t_{22}}t_{23} + 2t_{23} + 2qt_{11}v_2$ and (7) in $t_{33}| -\frac{(2v_1+1)t_{12}}{t_{11}t_{22}}t_{23} + qv_2 + 2qv_1v_2 + \frac{2v_1+1}{t_{11}}t_{13}$, but multiplying the second

term of the new condition (3) by $\frac{2v_1+1}{t_{11}}$ we obtain the new condition (7). Then the new set of conditions is

$$t_{33}|qt_{11}t_{22}, \quad t_{22}|t_{12}, \quad t_{33}| - \frac{2t_{12}}{t_{22}}t_{23} + 2t_{23} + 2qt_{11}v_2 \text{ and } t_{11}|2v_1 + 1.$$

Using Remark 2.2 we obtain that the integral (2.32) is

$$\begin{aligned} \zeta_{E,N,p}^{\leq}(s) &= (1 - p^{-1})^{-3} \int_{t_{33}|qt_{11}t_{22}} |t_{11}|^{s-1} |t_{22}|^{s-2} |t_{33}|^{s-3} d\mu \\ &= \zeta_p(s-2) (\zeta_p(s) \zeta_p(s-1) - p^{(2-s)(v_p(q)+1)} \zeta_p(2s-2) \zeta_p(2s-3)) \end{aligned}$$

If $p = 2$ then condition (5) says that t_{11} must be a unit, (6) is equivalent to $t_{22}|t_{12}$ and these conditions imply (2) and they also reduce (1) in $t_{33}|2qt_{22}$. This new condition (1) says that (4) is equivalent to $t_{33}|qt_{22}$ and therefore (1) is redundant, (3) can be reduced in $t_{33}| - \frac{2t_{12}}{t_{22}}t_{23} + 2t_{13} + 2qt_{11}v_2$ and (7) can be reduced in $t_{33}| - \frac{(2v_1+1)t_{12}}{t_{11}t_{22}}t_{23} + qv_2 + 2qv_1v_2 + \frac{2v_1+1}{t_{11}}t_{13}$ which is equivalent to $t_{33}| - \frac{t_{12}}{t_{22}}t_{23} + qt_{11} + qt_{11}v_2 + t_{13}$ because $\frac{v_1+1}{t_{11}}$ is a unit. Clearly this new condition (7) implies the new condition (3). Therefore the final set of conditions is:

$$t_{11} \in \mathbb{Z}_2^*, \quad t_{33}|qt_{22}, \quad t_{22}|t_{12}, \quad t_{33}| - \frac{t_{12}}{t_{22}}t_{23} + qt_{11} + qt_{11}v_2 + t_{13}.$$

The integral under this set of conditions is

$$\begin{aligned} \zeta_{E,N,2}^{\leq}(s) &= (1 - 2^{-1})^{-3} \int_{t_{33}|qt_{22}, t_{11} \in \mathbb{Z}_2^*} |t_{22}|^{s-2} |t_{33}|^{s-3} d\mu \\ &= \zeta_2(s-2) (\zeta_2(s-1) - 2^{-(s-2)(v_2(q)+1)} \zeta_2(2s-3)) \end{aligned}$$

Doing the product of all local factors we obtain:

$$\begin{aligned} \zeta_{E,N}^{\leq}(s) &= \zeta_2(s-2) (\zeta_2(s-1) - 2^{-(s-2)(v_2(q)+1)} \zeta_2(2s-3)) \cdot \\ &\quad \cdot \prod_{p \neq 2, p|q} \frac{\zeta_p(s) \zeta_p(s-1) - p^{(2-s)(v_p(q)+1)} \zeta_p(2s-2) \zeta_p(2s-3)}{\zeta_p(s) \zeta_p(s-1) - p^{2-s} \zeta_p(2s-2) \zeta_p(2s-3)} \\ &\quad \cdot \frac{\zeta(s) \zeta(s-1) \zeta(2s-2) \zeta(2s-3)}{\zeta(3s-3)} \end{aligned}$$

which has abscissa of convergence 2. The functional equation for almost all primes is:

$$(2.33) \quad \zeta_{E,N,p}^{\leq}(s)|_{p \rightarrow p^{-1}} = (-1)^3 p^{-3s+3} \zeta_{E,N,p}^{\leq}(s).$$

2.3.1. *The AB-groups of type $Q = p2gg$.* These are the groups

$$\begin{aligned} E : \langle x_1, x_2, x_3, \gamma_1, \gamma_2 : [x_2, x_1] = x_3^{4q}, [x_3, x_1] = [x_3, x_2] = 1, \\ \gamma_1 x_1 = x_1^{-1} \gamma_1 x_3^{2q}, \gamma_1 x_2 = x_2^{-1} \gamma_1 x_3^{-2q}, \gamma_1 x_3 = x_3 \gamma_1, \\ \gamma_2 x_1 = x_1 \gamma_2, \gamma_2 x_2 = x_2^{-1} \gamma_2 x_3^{-2q}, \gamma_2 x_3 = x_3^{-1} \gamma_2, \\ \gamma_1^2 = x_3, \gamma_2^2 = x_1, \gamma_1 \gamma_2 = x_1^{-1} x_2^{-1} \gamma_2 \gamma_1 x_3^{-(2q+1)} \rangle, \quad q \in \mathbb{N}. \end{aligned}$$

The Fitting subgroup is $N = N_{4q} = \langle x_1, x_2, x_3 \rangle$ and we have

$$(2.34) \quad \zeta_{E,N,p}^{\leq}(s) = (1 - p^{-1})^{-3} \int_{\mathcal{T}'} |t_{11}|^{s-3} |t_{22}|^{s-4} |t_{33}|^{s-5} d\mu$$

where $\mathcal{T}'$ is the set of pairs $(\mathbf{t}, \mathbf{v}) \in \text{Tr}(3, \mathbb{Z}_p) \times M_{2 \times 3}(\mathbb{Z}_p)$ satisfying:

$$\begin{aligned} t_{33} |4qt_{11}t_{22}, \quad (\gamma_j \mathbf{x}^{\mathbf{v}_i})^{-1} \mathbf{x}^{\mathbf{t}_i} \gamma_j \mathbf{x}^{\mathbf{v}_j} \in B_{\mathbf{t}}, \text{ for } i = 1, 2, 3, j = 1, 2 \\ (\gamma_1 \mathbf{x}^{\mathbf{v}_1})^2, (\gamma_2 \mathbf{x}^{\mathbf{v}_2})^2, (\gamma_1 \mathbf{x}^{\mathbf{v}_1} \gamma_2 \mathbf{x}^{\mathbf{v}_2})^2 \in B_{\mathbf{t}}. \end{aligned}$$

These condition are translated in:

- $t_{33} | 4qt_{11}t_{22}$
- $x_3^{2t_{i3}+2q(t_{i1}-t_{i2})+4q(t_{i1}v_{12}-v_{11}t_{i2})} \in B_{\mathbf{t}}$ for $i = 1, 2, 3$
- $x_2^{-2t_{i2}} x_3^{-2t_{i3}-2qt_{i2}-4q(t_{i1}v_{22}+v_{21}t_{i2})} \in B_{\mathbf{t}}$
- $x_3^{2v_{13}+2q(v_{11}-v_{12})+1-4qv_{11}v_{12}} \in B_{\mathbf{t}}$
- $x_1^{2v_{21}+1} x_3^{-2qv_{22}-4qv_{21}v_{22}} \in B_{\mathbf{t}}$
- $x_2^{-2v_{12}+2v_{22}-1} x_3^{-2q(v_{11}+v_{21})+4q(-v_{12}+v_{22})-2q+4q(v_{11}+v_{21})(-v_{12}+v_{22})} \in B_{\mathbf{t}},$

whose associated cone conditions are:

- (1) $t_{33} | 4qt_{11}t_{22}$
- (2) $t_{33} | 2t_{13} + 2q(t_{11} - t_{12}) + 4q(t_{11}v_{12} - v_{11}t_{12} - t_{11}t_{12})$
- (3) $t_{33} | 2t_{23} - 2q(1 + v_{11})t_{22}$
- (4) $t_{22} | 2t_{12}$
- (5) $t_{33} | \frac{2t_{12}}{t_{22}}t_{23} - 2t_{13} - 2qt_{12} - 4q(t_{11}v_{22} + v_{21}t_{12})$
- (6) $t_{33} | -2qt_{22} - 4qv_{21}t_{22}$
- (7) $t_{33} | 2v_{13} + 2q(v_{11} - v_{12}) + 1 - 4qv_{11}v_{12}$
- (8) $t_{11} | 2v_{21} + 1$
- (9) $t_{22} | \frac{2v_{21}+1}{t_{11}}t_{12}$
- (10) $t_{33} | -\frac{(2v_{21}+1)t_{12}}{t_{11}t_{22}}t_{23} + \frac{2v_{21}+1}{t_{11}}t_{13} + 2q(\frac{2v_{12}+1}{t_{11}})(\frac{2v_{12}+1}{t_{11}} - 1)t_{11}t_{12} + 2qv_{22} + 4qv_{21}v_{22}$
- (11) $t_{22} | -2v_{12} + 2v_{22} - 1$
- (12) $t_{33} | -\frac{-2v_{12}+2v_{22}-1}{t_{22}}t_{23} + q(v_{11} + 1 + 2v_{21} + 1)(-2v_{12} + 2v_{22} - 1)$

Suppose that $p \neq 2$. Using (1), (4) and (8), condition (2) can be reduced in $t_{33} | 2t_{13} + 2q(t_{11} - t_{12}) + 4q(t_{11}v_{12} - v_{11}t_{12})$ and condition (5) in $t_{33} | \frac{2t_{12}}{t_{22}}t_{23} - 2t_{13} - 4qt_{11}v_{22}$. Subtracting new condition (2) from condition (3) multiplied by t_{12}/t_{22} we obtain $t_{33} | \frac{2t_{12}}{t_{22}}t_{23} - 2t_{13} - 2qt_{12} - 2qt_{11}(1 +$

$2v_{12}$). This reduces new condition (5) in $t_{33}|2qt_{11}(-1+2v_{12}+2v_{22})$ and this new condition can be deduced from conditions (11) and (1). This shows that (5) is redundant. Condition (6) is also a consequence from (8) and (1), and (9) follows from (4). Conditions (4) and (1) reduce (10) in $t_{33}|-\frac{(2v_{21}+1)t_{12}}{t_{11}t_{22}}t_{23}+\frac{2v_{21}+1}{t_{11}}t_{13}+2qv_{22}+4qv_{21}v_{22}$, and this condition is implied by $t_{33}|-\frac{t_{12}}{t_{22}}t_{23}+t_{13}+2qt_{11}v_{22}$, and we saw that this is a consequence of condition (5). Finally condition (12) can be obtained from (3) multiplied by $\frac{-2v_{12}+2v_{22}-1}{t_{22}}$, in combination with (8) and (11). Then the final set of conditions is

- (1') $t_{33}|4qt_{11}t_{22}$
- (2') $t_{33}|2t_{13}+2q(t_{11}-t_{12})+4q(t_{11}v_{12}-v_{11}t_{12})$
- (3') $t_{33}|2t_{23}-2q(1+v_{11})t_{22}$
- (4') $t_{22}|2t_{12}$
- (5') $t_{33}|2v_{13}+2q(v_{11}-v_{12})+1-4qv_{11}v_{12}$
- (6') $t_{11}|2v_{21}+1$
- (7') $t_{22}| -v_{12}+2v_{22}-1$

Under these conditions the integral (2.34) is:

$$\begin{aligned}\zeta_{E,N,p}^{\leq}(s) &= (1-p^{-1})^{-3} \int_{t_{33}|qt_{11}t_{22}} |t_{11}|^{s-2}|t_{22}|^{s-2}|t_{33}|^{s-2} d\mu \\ &= \zeta_p(s-1)(\zeta_p(s-1)^2 - p^{-(s-1)(v_p(q)+1)}\zeta_p(2s-2)^2).\end{aligned}$$

If $p=2$, then conditions (7), (8) and (11) imply that t_{11}, t_{22} and t_{33} are units and then all the conditions are trivially satisfied. Then the integral (2.34) in that case is just

$$\zeta_{E,N,2}^{\leq}(s) = (1-2^{-1})^{-3} \int_{t_{11}, t_{22}, t_{33} \in \mathbb{Z}_2^*} d\mu = 1$$

Doing the product of all the local factors we obtain:

$$\begin{aligned}\zeta_{E,N}^{\leq}(s) &= \prod_{p \neq 2, p|q} \frac{\zeta_p(s-1)^2 - p^{-(s-1)(v_p(q)+1)}\zeta_p(2s-2)^2}{\zeta_p(s-1)^2 - p^{-(s-1)}\zeta_p(2s-2)^2} \\ &\quad \cdot \frac{\zeta(s-1)^2\zeta(2s-2)^2}{\zeta(3s-3)}\end{aligned}$$

which has abscissa of convergence 2. The functional equation for almost all primes is

$$\zeta_{E,N,p}^{\leq}(s)|_{p \rightarrow p^{-1}} = (-1)^3 p^{-3s+3} \zeta_{E,N,p}^{\leq}(s).$$

2.4. The AB-groups of type $Q = p4$. These are the groups

$$E : \langle x_1, x_2, x_3, \gamma : [x_2, x_1] = x_3^{2q}, [x_3, x_1] = [x_3, x_2] = 1 \\ \gamma x_1 = x_2 \gamma, \gamma x_2 = x_1^{-1} \gamma, \gamma^4 = x_3 \rangle, \quad q \in \mathbb{N}$$

and

$$F : \langle x_1, x_2, x_3, \gamma : [x_2, x_1] = x_3^{4q}, [x_3, x_1] = [x_3, x_2] = 1 \\ \gamma x_1 = x_2 \gamma, \gamma x_2 = x_1^{-1} \gamma, \gamma^4 = x_3^3 \rangle, \quad q \in \mathbb{N}$$

The Fitting subgroup of E is $N = N_{2q} = \langle x_1, x_2, x_3 \rangle$ while the Fitting subgroup for F is N_{4q} . Let G denote either E or F and let $\epsilon = 2$ or 4 according to when $G = E$ or F . We have

$$(2.35) \quad \zeta_{G,N,p}^{\leq}(s) = \int_{\mathcal{T}'} |t_{11}|^{s-2} |t_{22}|^{s-3} |t_{33}|^{s-4} d\mu$$

where $\mathcal{T}'$ is the set of pairs $(\mathbf{t}, \mathbf{v}) \in Tr(3, \mathbb{Z}_p) \times M_{1 \times 3}(\mathbb{Z}_p)$ such that

$$t_{33} | \epsilon q t_{11} t_{22}, \quad (\gamma \mathbf{x}^{\mathbf{v}})^{-1} \mathbf{x}^{\mathbf{t}_i} \gamma \mathbf{x}^{\mathbf{v}} \in B_{\mathbf{t}}, \text{ for } i = 1, 2, 3, \text{ and } (\gamma \mathbf{x}^{\mathbf{v}})^4 \in B_{\mathbf{t}}.$$

These conditions are translated in:

- $t_{33} | \epsilon q t_{11} t_{22}$
- $x_1^{t_{i2}} x_2^{-t_{i1}} x_3^{t_{i3} - \epsilon q t_{i2} v_2 - \epsilon q t_{12} t_{11} - \epsilon q v_{11} v_2 - \epsilon q v_1 t_{11}} \in B_{\mathbf{t}}, \text{ for } i = 1, 2, 3.$
- $x_3^{\epsilon - 1 + 4v_3 - 2\epsilon q v_1 v_2 - \epsilon q v_1^2 - \epsilon q v_2^2}.$

whose associated cone conditions are:

- (1) $t_{33} | \epsilon q t_{11} t_{22}$
- (2) $t_{11} | t_{12}$
- (3) $t_{22} | -\frac{t_{12}}{t_{11}} t_{12} - t_{11}$
- (4) $t_{33} | \frac{t_{11}^2 + t_{12}^2}{t_{11} t_{22}} t_{23} + (1 - \frac{t_{12}}{t_{11}}) t_{13} + \epsilon q \frac{1}{2} \frac{t_{12}}{t_{11}} (\frac{t_{12}}{t_{11}} + 1) t_{11} t_{12} + t_{13} - \epsilon q t_{12} v_2 - \epsilon q v_1 v_2 - \epsilon q v_1 t_{11}$
- (5) $t_{11} | t_{22}$
- (6) $t_{22} | -\frac{t_{22}}{t_{11}} t_{12}$
- (7) $t_{33} | (1 + \frac{t_{12}}{t_{11}}) t_{23} - \frac{t_{22}}{t_{11}} t_{13} + \epsilon q \frac{1}{2} \frac{t_{22}}{t_{11}} (\frac{t_{22}}{t_{11}} + 1) t_{11} t_{12} - \epsilon q v_2 t_{22} - \epsilon q v_1 v_2$
- (8) $t_{33} | \epsilon - 1 + 4v_3 - 2\epsilon q v_1 v_2 - \epsilon q v_1^2 - \epsilon q v_2^2$

Suppose that $p \neq 2$. We shall consider two cases:

Case 1: $t_{22} | t_{12}$. In this case conditions (2), (3), (5) and (6) are equivalent to $|t_{11}| = |t_{22}| \geq |t_{12}|$. Multiplying condition (7) by $\frac{t_{11} - t_{12}}{t_{22}}$ we can reduce condition (4) in $t_{33} | 2 \frac{t_{11}}{t_{22}} t_{23} + u$ where u is a function in the other variables which doesn't involve t_{23} nor t_{13} . Observe that $2 \frac{t_{11}}{t_{22}}$, which is the coefficient of t_{23} in (4), is a unit and the same is true for the coefficient of t_{13} in (7). Then we can compute the integral (2.35)

according to Remark 2.2. The result is:

$$\begin{aligned} & (1 - p^{-1})^{-3} \int_{\substack{t_{33}|qt_{11}t_{22}, \\ |t_{11}|=|t_{22}|}} |t_{11}|^{s-2} |t_{22}|^{s-2} |t_{33}|^{s-1} d\mu = \\ & = \zeta_p(s) (\zeta_p(2s-2) - p^{-s(v_p(q)+1)} \zeta_p(4s-2)) \end{aligned}$$

Case 2: $t_{22} \nmid t_{12}$. In this case conditions (2), (3), (5) and (6) become equivalent to condition (3) plus $|t_{11}| = |t_{12}| > |t_{22}|$. If we write $t_{12} = (k + pt)t_{11}$ for $k \in \{1, \dots, p-1\}$, then condition (3) implies $t_{22} | ((k + pt)^2 + 1)t_{11}$ and since $|t_{22}| < |t_{11}|$ then condition (3) implies $k^2 + 1 \equiv 0 \pmod{p}$. Since the equation $x^2 + 1 \equiv 0 \pmod{p}$ has solution $\pmod{p}$ if and only if $p \equiv 1 \pmod{4}$, then it makes sense to consider only the case $p \equiv 1 \pmod{4}$. In this case (by Hensel's Lemma) there exists $i \in \mathbb{Z}_p$ such that $i^2 + 1 = 0$. Then condition (3) is equivalent to $\frac{t_{22}}{t_{11}} | (\frac{t_{12}}{t_{11}} - i)(\frac{t_{12}}{t_{11}} + i)$ and this can happen if and only if $\frac{t_{22}}{t_{11}} | (\frac{t_{12}}{t_{11}} - i)$ or $\frac{t_{22}}{t_{11}} | (\frac{t_{12}}{t_{11}} + i)$ and both cases cannot occur at the same time. Then once t_{11} and t_{22} have been chosen, t_{12} can be chosen in a set of measure $2|t_{22}|$. For any such t_{11}, t_{22} and t_{12} then we have that $\frac{t_{11}+t_{12}}{t_{11}}$ and $\frac{t_{12}-t_{11}}{t_{11}}$ are both units. Multiplying condition (4) by $\frac{t_{22}}{t_{12}-t_{11}}$ we can eliminate the term which contains t_{13} in (7) and the coefficient of t_{23} in (7) will be $1 + \frac{t_{12}}{t_{11}} - \frac{1}{t_{12}-t_{11}} \frac{t_{11}^2+t_{12}^2}{t_{11}} = 1 - \frac{t_{12}+t_{11}}{t_{12}-t_{11}} = -2 \frac{t_{11}}{t_{12}-t_{11}}$ which is a unit. As in the other case, the integral we have to compute is

$$\begin{aligned} & 2 \cdot (1 - p^{-1})^{-3} \int_{\substack{t_{33}|qt_{11}t_{22} \\ |t_{11}|>|t_{22}|}} |t_{11}|^{s-2} |t_{22}|^{s-2} |t_{33}|^{s-1} d\mu = \\ & = \zeta_p(s) p^{-(s-1)} (\zeta_p(s-1) \zeta_p(2s-2) - p^{-s(v_p(q)+2)} \zeta_p(4s-2) \zeta_p(2s-1)) \end{aligned}$$

Adding the results in both cases we obtain that for $p \neq 2$:

$$\zeta_{G, N_{eq}, p}^{\leq}(s) = \zeta_p(s) (\zeta_p(2s-2) - p^{-s(v_p(q)+1)} \zeta_p(4s-2))$$

$$+ (\chi_4(p)+1) \zeta_p(s) p^{-(s-1)} (\zeta_p(s-1) \zeta_p(2s-2) - p^{-s(v_p(q)+2)} \zeta_p(4s-2) \zeta_p(2s-1)).$$

Suppose that $p = 2$. In that case condition (8) says that t_{33} must be a unit. Assuming that t_{33} is unit then conditions (4), (7) and (8) are trivially satisfied. We consider two cases:

Case 1: $t_{22} | t_{12}$. In this case (2), (3), (5) and (6) are equivalent to $|t_{11}| = |t_{22}| \geq |t_{12}|$ and then the integral is just

$$\begin{aligned} & (1 - 2^{-1})^{-3} \int_{\substack{t_{33} \in \mathbb{Z}_2^* \\ |t_{11}|=|t_{22}|}} |t_{11}|^{s-2} |t_{22}|^{s-2} d\mu = \\ & = \zeta_2(2s-2). \end{aligned}$$

Case 2: $t_{22} \nmid t_{12}$. In this case we have $|t_{11}| = |t_{12}| > |t_{22}|$. If $t_{12} = (1 + 2t)t_{11}$ then condition (2) is $t_{22}|(2 + 4t + 4t^2)t_{11}$ which means that $|t_{22}| = 2^{-1}|t_{11}|$. Then the integral is just

$$(1 - 2^{-1})^{-3} \int_{\substack{t_{33} \in \mathbb{Z}_2^* \\ |t_{11}|=2|t_{22}|}} |t_{11}|^{s-2} |t_{22}|^{s-2} d\mu = 2^{-(s-1)} \zeta_2(2s-2)$$

Summing the results in both cases we obtain

$$\zeta_{G, N_{eq}, 2}^{\leq}(s) = (1 + 2^{-(s-1)}) \zeta_2(2s-2) = \zeta_2(s-1).$$

Doing the product of all the local factors we obtain:

$$\begin{aligned} \zeta_{G, N_{eq}}^{\leq}(s) &= \zeta_2(s-1) \prod_{p \neq 2} [\zeta_p(s)(\zeta_p(2s-2) - p^{-s(v_p(q)+1)} \zeta_p(4s-2)) \\ &\quad + (\chi_4(p) + 1) \zeta_p(s) p^{-(s-1)} (\zeta_p(s-1) \zeta_p(2s-2) - p^{-s(v_p(q)+2)} \zeta_p(4s-2) \zeta_p(2s-1))] \\ &\quad \cdot \prod_{p|2q} \frac{L(3s-2, \chi_4, p)}{\zeta_p(s-1) \zeta_p(2s-1) L(s-1, \chi_4, p) L(2s-1, \chi_4, p)} \\ &\quad \cdot \frac{\zeta(s-1) \zeta(2s-1) L(s-1, \chi_4) L(2s-1, \chi_4)}{L(3s-2, \chi_4)} \end{aligned}$$

which has abscissa of convergence 2. The functional equation for almost all primes is:

$$(2.36) \quad \zeta_{G, N, p}^{\leq}(s)|_{p \rightarrow p^{-1}} = (-1)^3 \chi_4(p) p^{-3s+2} \zeta_{G, N}^{\leq}(s).$$

2.4.1. *The Bieberbach groups of type $Q = p3$.* These are the groups

$$E : \langle x_1, x_2, x_3, \gamma : [x_2, x_1] = x_3^{3q}, [x_3, x_1] = [x_3, x_2] = 1$$

$$\gamma x_1 = x_2 \gamma, \gamma x_2 = x_1^{-1} x_2^{-1} \gamma, \gamma x_3 = x_3 \gamma, \gamma^3 = x_3 \rangle, \quad q \in \mathbb{N},$$

$$F : \langle x_1, x_2, x_3, \gamma : [x_2, x_1] = x_3^{3q}, [x_3, x_1] = [x_3, x_2] = 1$$

$$\gamma x_1 = x_2 \gamma, \gamma x_2 = x_1^{-1} x_2^{-1} \gamma, \gamma x_3 = x_3 \gamma, \gamma^3 = x_3^2 \rangle, \quad q \in \mathbb{N},$$

$$G : \langle x_1, x_2, x_3, \gamma : [x_2, x_1] = x_3^r, [x_3, x_1] = [x_3, x_2] = 1$$

$$\gamma x_1 = x_2 \gamma x_3, \gamma x_2 = x_1^{-1} x_2^{-1} \gamma, \gamma x_3 = x_3 \gamma, \gamma^3 = x_3 \rangle, \quad r \in \mathbb{N}, r \notin 3\mathbb{N}.$$

Their Fitting subgroups are respectively N_{3q} , N_{3q} and N_r . Let G be any of the groups E, F and G , N its the Fitting subgroup and let k denote either $3q$ or r according to the case. Let also $\epsilon = 1$ if $G = E$ or G and $\epsilon = 2$ if $G = F$ and let $\delta = 1$ if $G = G$ and $\delta = 0$ in the other cases. We have

$$(2.37) \quad \zeta_{G, N, p}^{\leq}(s) = (1 - p^{-1})^{-3} \int_{\mathcal{T}'} |t_{11}|^{s-2} |t_{22}|^{s-3} |t_{33}|^{s-4} d\mu$$

where $\mathcal{T}'$ is the set of pairs $(\mathbf{t}, \mathbf{v}) \in Tr(3, \mathbb{Z}_p) \times M_{1 \times 3}(\mathbb{Z}_p)$ such that

- $t_{33} | kt_{11}t_{22}$
- $(\gamma^{\mathbf{x}^{\mathbf{v}_1}})^{-1} \mathbf{x}^{\mathbf{t}_i} \gamma^{\mathbf{x}^{\mathbf{v}_1}} \in B_{\mathbf{t}}$ for $i = 1, 2, 3$.
- $(\gamma^3 \mathbf{x}^{\mathbf{v}_1})^3 \in B_{\mathbf{t}}$.

These conditions are translated in:

- $t_{33} | kt_{11}t_{22}$
- $x_1^{-t_{11}+t_{12}} x_2^{-t_{11}} x_3^{t_{13}+kt_{11}\frac{t_{11}-1}{2}-kt_{11}t_{12}+kv_2t_{11}-kv_2t_{12}-kt_{11}v_1+\delta(t_{11}-t_{12})} \in B_{\mathbf{t}}$
- $x_1^{t_{22}} x_3^{t_{23}-kv_2t_{22}-\delta t_{22}} \in B_{\mathbf{t}}$
- $x_3^{3v_3-kv_1v_2-kv_2\frac{v_2+1}{2}-kv_1\frac{v_1+1}{2}+\epsilon+\delta(2v_1-v_2)} \in B_{\mathbf{t}},$

whose associated cone conditions are:

- (1) $t_{33} | kt_{11}t_{22}$
- (2) $t_{11} | -t_{11} + t_{12}$
- (3) $t_{22} | \frac{t_{11}^2 - t_{11}t_{12} + t_{12}^2}{t_{11}}$
- (4) $t_{33} | \frac{t_{11}^2 - t_{11}t_{12} + t_{12}^2}{t_{11}t_{22}} t_{23} + \frac{2t_{11}-t_{12}}{t_{11}} t_{13} + \frac{k}{2} \frac{-t_{11}+t_{12}}{t_{11}} \left(\frac{-t_{11}+t_{12}}{t_{11}} + 1 \right) t_{11}t_{12} + k(-t_{11} + t_{12})t_{11} + kt_{11}\frac{t_{11}-1}{2} - kt_{11}t_{12} + kv_2t_{11} - kv_2t_{12} - kt_{11}v_1 + \delta(t_{11} - t_{12}).$
- (5) $t_{11} | t_{22}$
- (6) $t_{22} | -\frac{t_{22}}{t_{11}} t_{12}$
- (7) $t_{33} | \frac{t_{11}+t_{12}}{t_{11}} t_{23} - \frac{t_{22}}{t_{11}} t_{13} + \frac{k}{2} \frac{t_{22}}{t_{11}} \left(\frac{t_{22}}{t_{11}} + 1 \right) t_{11}t_{12} - kv_2t_{22} - \delta t_{22}$
- (8) $t_{33} | 3v_3 - kv_1v_2 - kv_2\frac{v_2+1}{2} - kv_1\frac{v_1+1}{2} - kv_1v_2 + \epsilon + \delta(2v_1 - v_2)$

Suppose that $p \neq 3$. We consider two cases:

Case 1: $t_{22} | t_{12}$. In this case, condition (2), (3), (5) and (6) become $|t_{11}| = |t_{22}| \geq |t_{12}|$. Multiplying condition (7) by $\frac{2t_{11}-t_{12}}{t_{22}}$ and adding it to (3), we transform condition (3) in a new condition which doesn't involve t_{13} and where t_{23} has coefficient $3\frac{t_{11}}{t_{22}}$ which is a unit. Then we can compute the integral (2.37) under these conditions using Remak 2.2:

$$\begin{aligned} & (1-p^{-1})^{-3} \int_{\substack{t_{33} | kt_{11}t_{22} \\ |t_{11}| = |t_{22}|}} |t_{11}|^{s-2} |t_{22}|^{s-2} |t_{33}|^{s-1} d\mu \\ &= \zeta_p(s) (\zeta_p(2s-2) - p^{-s(v_p(k)+1)} \zeta_p(4s-2)) \end{aligned}$$

Case 2: $t_{22} \nmid t_{12}$. In this case we have that conditions (2), (3), (5) and (6) are equivalent to condition (3) plus $|t_{11}| = |t_{12}| > |t_{22}|$. Writing $t_{12} = (a+pt)t_{11}$ for $a \in \{1, \dots, p-1\}$ we see that condition (3) is equivalent to $t_{22} | t_{11}(1-a+a^2-pt+2apt+p^2t^2)$ which implies that $p | 1-a+a^2$. Since the equation $x^2-x+1=0$ has solution $\pmod{p}$ if and only if $p \equiv 1 \pmod{3}$ then it makes sense to consider only the case $p \equiv 1 \pmod{3}$. By Hensel's Lemma there exist $\alpha, \beta \in \mathbb{Z}_p^*$ such that $x^2-x+1 = (x-\alpha)(x-\beta)$. Then (3) is equivalent to $\frac{t_{22}}{t_{11}} | (\frac{t_{12}}{t_{11}} - \alpha)(\frac{t_{12}}{t_{11}} - \beta)$

and this can happen if and only if $\frac{t_{22}}{t_{11}}|\frac{t_{12}}{t_{11}} - \alpha$ or $\frac{t_{22}}{t_{11}}|\frac{t_{12}}{t_{11}} - \beta$ and these cases split the domain of integration in two disjoint sets. Thus, once t_{11} and t_{12} have been chosen we can choose t_{12} in a set of measure $|t_{22}|$. For t_{11}, t_{12} and t_{22} satisfying the conditions above we see clearly that $\frac{t_{11}-t_{12}}{t_{11}}$, which is the coefficient of t_{13} in (4), is a unit. Multiplying (4) by $\frac{t_{22}}{2t_{11}-t_{12}}$ (observe that $|2t_{11} - t_{12}| = |t_{11}|$, otherwise (3) cannot be satisfied) and adding it to condition (7) then we can transform (7) in a new condition which doesn't involve t_{13} and where the coefficient of t_{23} is $\frac{t_{11}^2-t_{11}t_{12}+t_{12}^2}{t_{11}t_{22}}\frac{t_{22}}{2t_{11}-t_{12}} + \frac{t_{11}+t_{12}}{t_{11}} = \frac{3t_{11}}{2t_{11}-t_{12}}$ which is a unit. Then we can compute the integral according to Remark 2.2. The result is

$$\begin{aligned} & 2(1-p^{-1})^{-3} \int_{\substack{t_{33}|kt_{11}t_{22} \\ |t_{11}|>|t_{22}|}} |t_{11}|^{s-2}|t_{22}|^{s-2}|t_{33}|^{s-1} d\mu \\ &= 2\zeta_p(s)p^{-(s-1)}(\zeta_p(s-1)\zeta_p(2s-2) - p^{-s(v_p(q)+2)}\zeta_p(4s-2)\zeta_p(2s-1)) \end{aligned}$$

Summing both cases we obtain for $p \neq 3$:

$$\begin{aligned} \zeta_{G,N,p}^{\leq}(s) &= \zeta_p(s)(\zeta_p(2s-2) - p^{-s(v_p(q)+1)}\zeta_p(4s-2)) + \\ & (1+\chi_3(p))\zeta_p(s)p^{-(s-1)}(\zeta_p(s-1)\zeta_p(2s-2) - p^{-s(v_p(q)+2)}\zeta_p(4s-2)\zeta_p(2s-1)). \end{aligned}$$

If $p = 3$ then for $k = 3q$ condition (8) says that t_{33} is a unit and for $k = r$, analyzing the residues mod 3 of r , v_{11} and v_{12} , we obtain that $3v_3 - rv_1v_2 - rv_2\frac{v_2+1}{2} - rv_1\frac{v_1+1}{2} + 1 + 2v_1 - v_2$ is not divisible by 3. Then we can assume that $t_{33} \in \mathbb{Z}_3^*$ in any case and therefore conditions (4), (7) and (8) are trivially satisfied. We consider two cases:

Case 1: $t_{22}|t_{12}$. The conditions are equivalent to $|t_{11}| = |t_{22}| \geq |t_{12}|$ and the integral is:

$$\begin{aligned} & (1-3^{-1})^{-3} \int_{\substack{t_{33} \in \mathbb{Z}_3^*, \\ |t_{11}|=|t_{22}|}} |t_{11}|^{s-2}|t_{s-2}|^{s-2} d\mu \\ &= \zeta_3(2s-2) \end{aligned}$$

Case 2: $t_{22} \nmid t_{12}$. In this case we have that the conditions are equivalent to condition (3) plus $|t_{11}| = |t_{12}| > |t_{22}|$. The solution of $x^2 - x + 1 \pmod 3$ is $x = 2$. Then t_{12} must be of the form $t_{12} = (2 + 3t)$ and condition (3) is therefore $t_{22}|(1-2+4-3t+12t+9t^2)t_{11} = (3+9t+9t^2)t_{11}$, which means that $|t_{22}| = 3^{-1}|t_{11}|$ and, as in the case of the groups of type $Q = p4$, this implies that t_{12} lives in a set of measure $|t_{22}|$. Then the integral is

$$\begin{aligned} & (1-3^{-1})^{-3} \int_{|t_{11}|=3|t_{22}|, t_{33} \in \mathbb{Z}_3^*} |t_{11}|^{s-2}|t_{22}|^{s-2} d\mu \\ &= 3^{-(s-1)}\zeta_3(2s-2) \end{aligned}$$

Summing the results in both cases we obtain

$$\zeta_{G,N,3}^{\leq}(s) = (1 + 3^{-(s-1)})\zeta_3(2s-2) = \zeta_3(s-1).$$

Doing the product of all the local factors we obtain:

$$\begin{aligned} \zeta_{G,N}^{\leq}(s) &= \zeta_3(s-1) \prod_{p \neq 3, p|k} \zeta_p(s) [(\zeta_p(2s-2) - p^{-s(v_p(k)+1)}\zeta_p(4s-2)) + \\ &(1 + \chi_3(p)) p^{-(s-1)}(\zeta_p(s-1)\zeta_p(2s-2) - p^{-s(v_p(k)+2)}\zeta_p(4s-2)\zeta_p(2s-1))] \\ &\cdot \prod_{p|k} \frac{L(3s-2, \chi_4, p)}{\zeta_p(s-1)\zeta_p(2s-1)L(s-1, \chi_3, p)L(2s-1, \chi_3, p)} \\ &\cdot \frac{\zeta(s-1)\zeta(2s-1)L(s-1, \chi_3)L(2s-1, \chi_3)}{L(3s-2, \chi_3)} \end{aligned}$$

which has abscissa of convergence 2. The functional equation for almost all primes is:

$$(2.38) \quad \zeta_{G,N,p}^{\leq}(s)|_{p \rightarrow p^{-1}} = (-1)^3 p^{-3s+2} \chi_4(p) \zeta_{G,N,p}^{\leq}(s).$$

2.4.2. *The Bieberbach groups of type $Q = p6$.* These are the groups:

$$\begin{aligned} E : \langle x_1, x_2, x_3, \gamma : [x_2, x_1] &= x_3^{6q}, [x_3, x_1] = [x_3, x_2] = 1 \\ \gamma x_1 &= x_1 x_2 \gamma, \gamma x_2 = x_1^{-1} \gamma, \gamma x_3 = x_3 \gamma, \gamma^6 = x_3 \rangle, \quad q \in \mathbb{N}, \end{aligned}$$

$$\begin{aligned} F : \langle x_1, x_2, x_3, \gamma : [x_2, x_1] &= x_3^{6q+4}, [x_3, x_1] = [x_3, x_2] = 1 \\ \gamma x_1 &= x_1 x_2 \gamma, \gamma x_2 = x_1^{-1} \gamma, \gamma x_3 = x_3 \gamma, \gamma^6 = x_3 \rangle, \quad q \in \mathbb{N}, \end{aligned}$$

$$\begin{aligned} G : \langle x_1, x_2, x_3, \gamma : [x_2, x_1] &= x_3^{6q}, [x_3, x_1] = [x_3, x_2] = 1 \\ \gamma x_1 &= x_1 x_2 \gamma, \gamma x_2 = x_1^{-1} \gamma, \gamma x_3 = x_3 \gamma, \gamma^6 = x_3^5 \rangle, \quad q \in \mathbb{N}, \end{aligned}$$

$$\begin{aligned} H : \langle x_1, x_2, x_3, \gamma : [x_2, x_1] &= x_3^{6q+2}, [x_3, x_1] = [x_3, x_2] = 1 \\ \gamma x_1 &= x_1 x_2 \gamma, \gamma x_2 = x_1^{-1} \gamma, \gamma x_3 = x_3 \gamma, \gamma^6 = x_3^5 \rangle, \quad q \in \mathbb{N}. \end{aligned}$$

Let G denote any of the groups E , F , G and H and let k denote $6q$, $6q+2$ or $6q+4$ according to the case. Let N be the Fitting subgroup of G and let $\epsilon = 1$ if G is any of the groups E , F and $\epsilon = 5$ in the other cases. We have:

$$(2.39) \quad \zeta_{G,N,p}^{\leq}(s) = (1 - p^{-1})^{-3} \int_{\mathcal{T}'} |t_{11}|^{s-2} |t_{22}|^{s-3} |t_{33}|^{s-3} d\mu$$

where $\mathcal{T}'$ is the set of pairs $(\mathbf{t}, \mathbf{v}) \in \text{Tr}(3, \mathbb{Z}_p) \times M_{1 \times 3}(\mathbb{Z}_p)$ such that

- $t_{33} | k t_{11} t_{22}$
- $(\gamma \mathbf{x}^{\mathbf{v}_1})^{-1} \mathbf{x}^{\mathbf{t}_i} \gamma \mathbf{x}^{\mathbf{v}_1} \in B_{\mathbf{t}}$ for $i = 1, 2, 3$.
- $(\gamma \mathbf{x}^{\mathbf{v}_1})^6 \in B_{\mathbf{t}}$.

These conditions are translated in:

- $t_{33} | kt_{11}t_{22}$
- $x_1^{t_{12}} x_2^{-t_{11}+t_{12}} x_3^{t_{13}+kt_{12}(t_{12}+1)-kt_{11}t_{12}-kv_2t_{12}-kt_{11}v_1+kt_{12}v_1} \in B_t$
- $x_1^{t_{22}} x_2^{t_{22}} x_3^{t_{23}+6qt_{22}(t_{22}+1)-6qv_2t_{22}+6qt_{22}v_1} \in B_t$
- $x_3^{1+6v_3-kv_1(3v_1+3)-kv_2(3v_2-3)} \in B_t.$

whose associated cone conditions are:

- (1) $t_{33} | kt_{11}t_{22}$
- (2) $t_{11} | t_{12}$
- (3) $t_{22} | -\frac{t_{12}^2+t_{11}^2-t_{11}t_{12}}{t_{11}}$
- (4) $t_{33} | \frac{t_{12}^2+t_{11}^2-t_{11}t_{12}}{t_{11}t_{22}}t_{23} + (1-\frac{t_{12}}{t_{11}})t_{13} + \frac{k}{2}\frac{t_{12}}{t_{11}}(\frac{t_{12}}{t_{11}}+1)t_{11}t_{12} - kt_{12}(-t_{11}+t_{12}) + kt_{12}(t_{12}+1) - kt_{11}t_{12} - kv_2t_{12} - kt_{11}v_1 + kt_{12}v_1$
- (5) $t_{11} | t_{22}$
- (6) $t_{22} | -\frac{t_{22}}{t_{11}}t_{12} + t_{22}$
- (7) $t_{33} | \frac{t_{12}}{t_{11}}t_{23} - \frac{t_{22}}{t_{11}}t_{13} + \frac{k}{2}\frac{t_{22}}{t_{11}}(\frac{t_{22}}{t_{11}}+1) - kt_{22}^2 + kt_{22}(t_{22}+1) - kv_2t_{22} + kt_{22}v_1$
- (8) $t_{33} | \epsilon + 6v_3 - kv_1(3v_1+3) - kv_2(3v_2-3).$

Suppose that $p \neq 2, 3$. We consider two cases:

Case 1: $t_{22} | t_{12}$. In this case, condition (2), (3), (5) and (6) are equivalent to $|t_{11}| = |t_{22}| \geq |t_{12}|$. Multiplying condition (7) by $\frac{t_{11}-t_{12}}{t_{22}}$ and adding it to condition (4) we can eliminate the term which contains t_{13} in (4) and the coefficient of t_{23} will be $\frac{t_{12}^2+t_{11}^2-t_{11}t_{12}}{t_{11}t_{22}} + \frac{t_{11}-t_{12}}{t_{22}} \frac{t_{12}}{t_{11}} = \frac{t_{11}}{t_{22}}$ which is a unit. Then we can compute the integral as we did in the last cases:

$$\begin{aligned} & (1-p^{-1})^{-3} \int_{\substack{t_{33} | kt_{11}t_{22} \\ |t_{11}|=|t_{22}|}} |t_{11}|^{s-2} |t_{22}|^{s-2} |t_{33}|^{s-1} d\mu \\ &= \zeta_p(s) (\zeta_p(2s-2) - p^{-s(v_p(k)+1)} \zeta_p(4s-2)) \end{aligned}$$

Case 2: $t_{22} \nmid t_{12}$. In this case conditions (2),(3),(5) and (6) are equivalent to condition (2) plus $|t_{11}| = |t_{12}| > |t_{22}|$. We saw in the case of groups of type $Q = p3$ that condition (3) can be satisfied if and only if $p \equiv 1 \pmod{3}$, and for $p \equiv 1 \pmod{3}$ we have that the set of those t_{12} satisfying this condition has measure $2|t_{22}|$. Then the integral is:

$$\begin{aligned} & 2(1-p^{-1})^{-3} \int_{\substack{t_{33} | qt_{11}t_{22} \\ |t_{11}|>|t_{22}|}} |t_{11}|^{s-2} |t_{22}|^{s-2} |t_{33}|^{s-1} d\mu \\ &= 2\zeta_p(s) p^{-(s-1)} (\zeta_p(s-1) \zeta_p(2s-2) - p^{-s(v_p(q)+2)} \zeta_p(4s-2) \zeta_p(2s-1)) \end{aligned}$$

Summing the results in both cases we obtain for $p \neq 2, 3$

$$\begin{aligned} \zeta_{G,N,p}^{\leq}(s) &= \zeta_p(s)(\zeta_p(2s-2) - p^{-s(v_p(q)+1)}\zeta_p(4s-2)) + \\ &+ (1 + \chi_3(p))\zeta_p(s)p^{-(s-1)}(\zeta_p(s-1)\zeta_p(2s-2) - p^{-s(v_p(q)+2)}\zeta_p(4s-2)\zeta_p(2s-1)). \end{aligned}$$

If $p = 2$ or $p = 3$ then condition (8) implies that t_{33} must be a unit and the integral is the same as that for groups E of type $Q = p3$ giving the same result except for the case $p = 2$ where we only have to consider the case $t_{22}|t_{12}$. The results are:

$$\zeta_{G,N,2}^{\leq}(s) = \zeta_2(2s-2) \text{ and } \zeta_{G,N,3}^{\leq}(s) = \zeta_3(s-1).$$

Doing the product of all the local factors we obtain:

$$\begin{aligned} \zeta_{G,N}^{\leq}(s) &= \zeta_2(2s-2)\zeta_3(s-1) \prod_{p \neq 2,3} \zeta_p(s) [(\zeta_p(2s-2) - p^{-s(v_p(q)+1)}\zeta_p(4s-2)) + \\ &(1 + \chi_3(p))p^{-(s-1)}(\zeta_p(s-1)\zeta_p(2s-2) - p^{-s(v_p(q)+2)}\zeta_p(4s-2)\zeta_p(2s-1))] \\ &\prod_{p \neq 2,3; p|k} \frac{L(3s-2, \chi_4, p)}{\zeta_p(s-1)\zeta_p(2s-1)L(s-1, \chi_3, p)L(2s-1, \chi_3, p)} \\ &\cdot \frac{\zeta(s-1)\zeta(2s-1)L(s-1, \chi_3)L(2s-1, \chi_3)}{L(3s-2, \chi_3)} \end{aligned}$$

which has abscissa of convergence 2. The functional equation for almost all primes is:

$$(2.40) \quad \zeta_{G,N,p}^{\leq}(s)|_{p \rightarrow p^{-1}} = (-1)^3 p^{-3s+2} \chi_3(p) \zeta_{G,N,p}^{\leq}(s).$$

ACKNOWLEDGEMENTS

The author's work was supported by CONICET and it was mainly developed during his visit to KULeuven-Kulak with an ERASMUS scholarship. The author is very grateful for the hospitality in KULeuven and in particular he thanks Karel Dekimpe for provided him with all the facilities to do this work. The author also thanks to the GAP System developers. It was very useful to check all the computations.

REFERENCES

- [D] K. Dekimpe, *Almost-Bieberbach Groups: Affine and Polynomial Structures*. Lecture Notes in Mathematics 1639. Springer-Verlag Berlin-Heidelberg 1996.
- [DIKL] K. Dekimpe, P. Igodt, S. Kim and K. Lee, *Affine structures for closed 3-dimensional manifolds with nil-geometry*. Quart. J. Math. Oxford Ser. (2) 46 (1995), no. 182, 141Ü167. 57S30 (57M50)
- [dSMS] M. P. F. du Sautoy, J. J. McDermott and G. C. Smith, *Zeta Functions of Crystallographic Groups and Analytic Continuation* Precedings of the London Mathematical Society, Vol. 79, Issue 3, 511-534.

- [dSG] M. P. F. du Sautoy, F. Grunewald, *Analytic properties of zeta functions and subgroup growth*. Ann. of Math. **152**(2) (2000), 793-833.
- [dSW] M. P. F. du Sautoy, L. Woodward, *Zeta functions of Groups and Rings* Lectures note in mathematics 1925, Springer-Verlag, 2008.
- [GSS] F. Grunewald, D. Segal, G. C. Smith, *Subgroups of finite index in nilpotent groups*. Invent. Math. **93** (1988), no. 1, 185-223.
- [LMS] A. Lubotzky, A. Mann and D. Segal, *Finitely generated groups of polynomial subgroup growth*. Israel J. Math. **82**(1993), 363-371.
- [S] D. Sulca, *Zeta functions of virtually nilpotent groups* Arxiv: <http://128.84.158.119/abs/1205.4464>.
- [V] C. Voll, *Functional equation of zeta functions of groups and rings*, Ann. of Math. **172**(2010), 1181-1218.

UNIVERSIDAD NACIONAL DE CÓRDOBA, CIUDAD UNIVERSITARIA, 5000 CÓRDOBA.

E-mail address: `sulca@famaf.unc.edu.ar`